

VALIDATION OF BIOLUMINESCENT *ESCHERICHIA COLI* O157:H7 FOR USE AS
A PRE-HARVEST FOOD SAFETY MODEL

By

Heather Ann Duoss

A Thesis
Submitted to the Faculty of
Mississippi State University
in Partial Fulfillment of the Requirements
for the Degree of Master of Science
in Agriculture and Life Sciences
in the Department of Animal and Dairy Sciences

Mississippi State, Mississippi

May 2012

Copyright 2012

By

Heather Ann Duoss

VALIDATION OF BIOLUMINESCENT *ESCHERICHIA COLI* O157:H7 FOR USE AS
A PRE-HARVEST FOOD SAFETY MODEL

By

Heather Ann Duoss

Approved:

Ty Schmidt
Assistant Professor of
Animal and Dairy Sciences
(Director of Thesis)

Todd Callaway
Adjunct Professor of
Animal and Dairy Sciences
(Committee Member)

Janet Donaldson
Assistant Professor of
Biological Sciences
(Committee Member)

James (Mike) Martin
Assistant Professor of
Animal and Dairy Sciences
(Committee Member)

Brian Rude
Professor of
Animal and Dairy Sciences
(Graduate Coordinator)

George Hopper
Dean of
College of Agriculture and Life Science

Name: Heather Ann Duoss

Date of Degree: May 11, 2012

Institution: Mississippi State University

Major Field: Agriculture and Life Sciences

Major Professor: Ty Schmidt

Title of Study: VALIDATION OF BIOLUMINESCENT *ESCHERICHIA COLI*
O157:H7 FOR USE AS A PRE-HARVEST FOOD SAFETY MODEL

Pages in Study: 95

Candidate for Degree of Master of Science

Cattle are naturally colonized by enterohemorrhagic *Escherichia coli* within the gastrointestinal tract. The most notorious of the enterohemorrhagic *E. coli* is *E. coli* O157:H7, which can cause serious illness to humans if ingested. To ensure that the United States has a safe food supply, research is ongoing in pre-harvest food safety and pathogen intervention strategies. While advances in pre-harvest intervention strategies are encouraging, no method has proven to completely eliminate and/or control O157:H7. A key limitation to successful pathogen intervention strategies is the inability to track and monitor pathogens in a real-time fashion. Through the use of bioluminescent plasmids harboring the *luxCDABE* cassette, pathogen tracking could be a viable solution. Bioluminescent plasmids are capable of facilitating the tracking, pathogenesis and physical locations of pathogens, thus enabling researchers to have a better understanding of the pathogenic process.

DEDICATION

I would like to dedicate this thesis and all the work required to obtain my Masters degree to my family and especially to the most important person in my life, my fiancé Zack. Thank you Dad and Mom for always being supportive in anything I have ever done and thank you for always giving me that extra push and love I needed most of the time. My little brother Derek, thank you for always being that quiet but yet supportive brother. Thank you to my little sister Jessi for always giving me something to laugh at when I needed it and someone to talk to when I needed it! Grandma and Grandpa, thank you for always being there for me and attending every event I have ever had, whether it is a meeting, sports game, graduation or anything else. I couldn't have done it without any of you. I would also like to thank my Aunt Carmen and Teagan for all the support you have given me in my life, I love you! Finally, I owe a special thank you to my fiancé Zack for all the pain and torture I have put you through during my Masters degree. You have been the rock that has helped me get through this. I appreciate everything you have done for me and everything you have put up with! I love you and thank you for everything.

ACKNOWLEDGEMENTS

I have a lot of people to thank for all the help I have received throughout my Masters degree. During my time here at Mississippi State University I have had the great pleasure of working with some extremely great people. I would like to thank the graduate and undergraduate students who have helped me along the way. Thank you to Katie Pfeiffer, Badger, for always being there and being a great friend. Landon Marks, Cheese, for always giving me something to laugh at and shake my head at. Also, thank you for never letting any of us miss lunch! Rand Broadway, for all the help and insight you have given me on my research projects. My academic big brother Joe Buntyn and big sister Shollie Falkenberg, thank you for your honesty and always looking out for me.

I would also like to thank Dr. Carolyn Huntington for allowing me to be a teaching assistant for Introduction to Animal Science. It was a great opportunity and I wouldn't trade it for the world. Thank you to all my undergraduate students who kept me on my toes and always, some way or another, made me laugh.

To my committee, Dr. Janet Donaldson, Dr. Todd Callaway, Dr. Mike Martin and Dr. Ty Schmidt thank you for your advice and guidance along the way. My research couldn't have been possible without all of your help. Dr. Donaldson, thank you for all the time and effort you have put into understanding the Animal Science language and techniques. I am sad to say that the rumen and fecal fluid is now in your hands! You have been a great help to me and you were always there whenever I needed anything. Dr. Callaway, thank you for all the knowledge and insight you have given me over the past

two years on all my projects. Your sense of humor is always needed and is always a good sigh of relief. Dr. Martin, thank you for always giving great advice and a history lesson, you are one of the most knowledgeable people I know. Finally, my major professor Dr. Schmidt, thank you for always being there for me and everything you have ever done for me. You have been a great professor, researcher and mentor. Thank you for taking me into your lab and giving me a chance to figure out what this graduate school thing is all about. You have given me many opportunities and helped me gain many acquaintances along the way, and I am truly thankful for that.

TABLE OF CONTENTS

	Page
DEDICATION	ii
ACKNOWLEDGEMENTS	iii
LIST OF TABLES	viii
LIST OF FIGURES	ix
CHAPTER	
I. INTRODUCTION	1
II. LITERATURE REVIEW	3
<i>Escherichia coli</i> O157:H7 and pre-harvest food safety	3
Growth and reproduction of <i>Escherichia coli</i>	4
<i>Escherichia coli</i> virulence groups	5
<i>Escherichia coli</i> O157:H7	7
Shiga toxin-producing <i>Escherichia coli</i> (STEC)	8
Role of enterohemorrhagic <i>Escherichia coli</i> in ruminant animals	11
<i>Escherichia coli</i> O157:H7 shedding in cattle	13
Prevalence of <i>Escherichia coli</i> O157:H7 on cattle hides	13
<i>Escherichia coli</i> pre-harvest intervention strategies	14
Management practices and transport	15
Cattle water and feed management	16
Live animal <i>Escherichia coli</i> O157:H7 treatments	20
Citrus by-products	22
Future of pre-harvest	24
Biophotonics and bioluminescence	25
Successful models	26
Advantages of bioluminescence	26
Literature cited	28
III. SURVIVAL OF <i>ESCHERICHIA COLI</i> O157:H7 (ATCC 43888) TRANSFORMED WITH EITHER THE PAK1- <i>LUX</i> OR PXEN-13 PLASMIDS IN BOVINE MIXED RUMINAL AND FECAL MICROORGANISM FLUID	48

Abstract.....	48
Introduction.....	49
Materials and Methods.....	51
Bacterial Cultivation Conditions.....	51
Pure Culture Growth.....	51
Mixed Ruminant and Fecal Microorganism Contents Collection.....	52
Mixed Ruminant and Fecal Microorganism Contents Fermentations.....	53
Statistical Analysis.....	53
Results.....	54
Growth of <i>E. coli</i> 43888 containing pAK1- <i>lux</i> or pXEN-13.....	54
Growth in Ruminant Fluid Fermentations.....	54
Growth in Fecal Fluid Fermentations.....	55
Comparison of Photonic Emissions by pAK1- <i>lux</i> and pXEN-13.....	55
Discussion.....	56
Conclusions.....	58
Literature cited.....	62
IV. EFFECTS OF CITRUS BY-PRODUCTS ON GROWTH OF O157:H7 AND NON-O157 <i>ESCHERICHIA COLI</i> SEROTYPES WITHIN BOVINE MIXED RUMINAL MICROORGANISM FLUID.....	65
Abstract.....	65
Introduction.....	66
Materials and Methods.....	68
Ruminant Fluid Collection and Medium Preparation.....	68
Bacterial Serotypes and Cultivation Conditions.....	68
Citrus By-Product Trial.....	69
Statistical Analysis.....	69
Results.....	70
Citrus By-Products pH Values.....	70
Powdered Citrus By-Product.....	70
Pelleted Citrus By-Product.....	71
Discussion.....	71
Conclusion.....	73
Literature cited.....	76
V. EVALUATION OF THE COLONIZATION OF <i>ESCHERICHIA</i> <i>COLI</i> O157:H7 WITHIN THE SWINE GASTROINTESTINALTRACT.....	79
Abstract.....	79
Introduction.....	80
Materials and Methods.....	81
Bacterial Cultivation Conditions.....	81
Swine Trial.....	82

Gastrointestinal Tract Sections	82
Gastrointestinal Tract Rinse.....	83
Statistical Analysis.....	83
Results.....	84
Gastrointestinal Tract Contents.....	84
Gastrointestinal Tract Rinse.....	85
Discussion.....	86
Conclusion	88
Literature cited	91
VI. CONCLUSION.....	94

LIST OF TABLES

TABLE	Page
4.1 Least squares means for growth of enterohemorrhagic <i>Escherichia coli</i> (EHEC) O26:H11, O103:H8, O111:H8, O145:H28, O157:H7 and O157:H7 <i>Dstx₁stx₂</i> within bovine mixed rumen microorganism fluid medium, supplemented with 0%, 5%, 10% and 20% powdered citrus by-product (CBP) over time (h)	74
4.2 Least squares means for growth of enterohemorrhagic <i>Escherichia coli</i> (EHEC) O26:H11, O103:H8, O111:H8, O145:H28, O157:H7 and O157:H7 <i>Dstx₁stx₂</i> within bovine mixed rumen microorganism fluid medium, supplemented with 0%, 5%, 10% and 20% pelleted citrus by-product (CBP) over time (h).....	75
5.1 The least square means of <i>Escherichia coli</i> O157:H7+pAK1- <i>lux</i> (ATCC 43888) populations (log ₁₀ CFU/mL) within contents collected from the gastrointestinal tract of swine	89
5.2 The least square means of <i>Escherichia coli</i> O157:H7+pAK1- <i>lux</i> (ATCC 43888) populations (log ₁₀ CFU/mL) within rinse obtained from the gastrointestinal tract of swine	90

LIST OF FIGURES

FIGURE	Page
3.1	The OD ₆₀₀ of cultures of the wild type (WT) <i>Escherichia coli</i> O157:H7 ATCC 43888 (◆) and the two bioluminescently transformed variants, 43888 containing the plasmid pXEN-13 (▲) or pAK1- <i>lux</i> (■). Cultures were grown in the absence of (A) and presence (B) of ampicilloin (50 mg/mL) to provide selective pressure.....59
3.2	The least square means of <i>Escherichia coli</i> O157:H7 (ATCC 43888) populations (log ₁₀ CFU/mL) in bovine mixed ruminal microorganism incubations. Growth was monitored by viable plate counts with the parental strain (◆; WT) or 43888 containing the plasmid pAK1- <i>lux</i> (▲; PAK) or pXEN-13 (■; XEN) in bovine mixed ruminal fluid for 6 h. Values represent the log ₁₀ CFU/mL values from three independent replicates.....60
3.3	The least square means of <i>Escherichia coli</i> O157:H7 (ATCC 43888) populations (log ₁₀ CFU/mL) in bovine mixed fecal microorganism incubations. Growth was monitored by viable plate counts with the parental strain (◆; WT) or 43888 containing the plasmid pAK1- <i>lux</i> (▲; PAK) or pXEN-13 (■; XEN) in bovine mixed fecal fluid for 6 h. Values represent the log ₁₀ CFU/mL values from three independent replicates.....61

CHAPTER I

INTRODUCTION

In 2011, *Escherichia coli* O157:H7 was responsible for the recalls of an estimated 1 million pounds of ground beef due to possible contamination (USDA-FSIS, 2011). *Escherichia coli* O157:H7 was first identified as a pathogen in 1982 (Wetzel and LeJeune 2006) and was linked to undercooked meat. The primary means for human contraction of *E. coli* O157:H7 is through the ingestion of raw or undercooked meat, water runoff from cattle environments (Armstrong et al., 1996) and/or animal-to-person contact (Wetzel and LeJeune, 2006).

Cattle are a reservoir for O157:H7 and contribute to the spread, as well as the epidemiology of human O157:H7 infection (Griffin and Tauxe, 1991; Sargeant et al., 2006). *Escherichia coli* O157:H7 has the ability to colonize within the gastrointestinal tract of cattle without inflicting clinical illness and subsequently enabling cattle to become a host for O157:H7 (Low et al., 2005). Cattle lack the globotriaosylceramide (Gb₃) receptor shiga toxins within the large intestine bind to, thus why cattle are able to harbor O157:H7 within the gastrointestinal tract but remain symptomatic (Hoey et al., 2002). *Escherichia coli* O157:H7 prevalence studies estimate that approximately 30% of cattle throughout the United States harbor this pathogen in their feces (Elder et al., 2000; Callaway et al., 2003).

Many pre-harvest strategies have been implemented by researchers to potentially reduce and/or eliminate O157:H7 within cattle such as vaccinations, direct-fed microbials

and diet manipulations. Although some intervention strategies have proven to reduce O157:H7 populations, none have completely eradicated O157:H7 or proven to be repeatable and consistent. One of the main limitations associated with pre-harvest intervention development is the inability to physically track O157:H7 in real-time.

A potential solution to the inability to monitor pathogens is through the implementation of bioluminescent plasmids. Bioluminescent plasmids that contain the *luxCDABE* operon encode enzymes responsible for synthesis of the luciferase (Meighen, 1993). These plasmids can then be introduced into pathogens, which can then be monitored through an intensified charged-couple device camera (Ryan et al., 2011). Employing bioluminescence *in vivo* can potentially facilitate tracking of pathogens (Contag et al., 1997), allowing for the indication of the physical location (Siragusa et al., 1999) and thus increasing researchers understanding of pathogenic processes (Contag et al., 1995) of O157:H7.

CHAPTER II

LITERATURE REVIEW

***Escherichia coli* O157:H7 and pre-harvest food safety**

The bacterium *Escherichia coli* (*E. coli*) is one of the most commonly isolated bacteria in throughout the meat and food industries (Matic et al., 1997). *Escherichia coli* are gram-negative, facultative anaerobic, rod-shaped bacteria that are non-sporulating with an optimal growth at 37°C. There are over 700 known serotypes of *E. coli* (Ray and Schaffer, 2011). Serotyping as a means of identifying various serovars of *E. coli* is achieved by serotyping by antigens. In example *E.coli* O157:H7 indicates that the O antigen encompasses the cell wall outer membrane or somatic antigen number 157, while the H antigen indicates the flagella antigen 7. The O antigen is part of the lipopolysaccharide present in the outer membrane of bacteria (Sheng et al., 2008).

While most *E.coli* are non-pathogenic, certain serotypes of *E. coli* can be pathogenic; these pathogens account for devastating foodborne related diseases throughout the world (Doyle, 1991; Hilborn et al., 1999; Vogt and Dippold, 2005). Nonpathogenic *E. coli* are cultured in the normal flora of warm blooded animal's gastrointestinal (GI) tracts (Rembacken et al., 1999; Guarner and Malagelada, 2003). Nonpathogenic *E. coli* can also prevent pathogenic *E. coli* from colonizing within the intestine by providing a protective barrier (Hudault et al., 2000; Reid et al., 2001). Nonpathogenic bacteria can provide colonization resistance such as a protective barrier,

in which bacteria can uphold an ecological balance and prevent pathogenic bacteria from colonizing through competitive exclusion (van der Waaij et al., 1971; Rolfe, 2000).

Growth and reproduction of *Escherichia coli*

Typical growth patterns of *E. coli* and other bacteria are measured in four different phases: lag phase, log phase, stationary phase and death phase (Roszak and Colwell, 1987). When bacteria are introduced to medium, adjustment to the surrounding environment and growth conditions will occur, thus representing lag phase (Baranyi, 2002). The log phase or also known as exponential phase is characterized by cell growth, where bacteria exhibit continual doubling bacterial populations (Jensen and Hammer, 1993) until the nutrients of the medium utilized are depleted and cellular waste products accumulate (Kolter et al., 1993). When the growth rate begins to decrease due to lack of nutrients, bacteria will enter stationary phase. Stationary phase is characterized by a stable and constant rate of bacterial growth that is equivalent to the rate of bacterial death (Delpy et al., 1956). After all nutrients have been depleted death phase emerges and bacteria will begin to die.

Bacteria such as *E. coli* reproduce by means of binary fission. The single bacterium will elongate during the growth phase, reaching twice its length, and binary fission will occur (Angert, 2005). The single cell bacterium's DNA replicates and segregates into two identical chromosomes (Donachie, 2001). After the chromosomal segregation within the bacterium the individual chromosomes will move to opposite ends of the bacterium (Donachie, 2001). This allows for cytokinesis, which involves the actual splitting of the bacterium into two identical daughter cells (Angert, 2005).

***Escherichia coli* virulence groups**

Pathogenic *E. coli* can be broken down into five groups, based on virulence attributes (Johnson, 1991). These five groups are enterotoxigenic (ETEC), enteroaggregative (EA_ggEC), enteropathogenic (EPEC), enteroinvasive (EIEC) and enterohemorrhagic (EHEC; Levine, 1987). The ETEC utilize fimbriae, which are hair-like structures on the outer membrane and are shorter than flagella. The ETEC use fimbriae to attach to the specific receptors on enterocytes within the intestinal lumen (Levine and Edelman, 1984). The ETEC virulence group can produce heat-stable and/or heat labile enterotoxins (Moseley et al., 1980). This virulence group is the primary cause of traveler's diarrhea which results from an accumulation of fluid and electrolytes within the GI tract lumen (Levine, 1987). Common ETEC serogroups include O8, O139, O141, O147 and O149 (Nagy and Fekete, 1999).

Enteroaggregative *E. coli* (EA_ggEC) or also known as enteroadherent, are one of the newest *E. coli* virulence groups. The EA_ggEC have fimbriae similar to ETEC that attach to the intestinal epithelial cells and some strains are capable of producing a heat-stable enterotoxin (Kaper et al., 2004). The result from an EA_ggEC infection results in watery diarrhea, and is usually observed in children in developing countries (Nataro et al., 1995). In comparison to the other virulence groups, EA_ggEC has a longer incubation period within humans, which has been estimated to be 20 to 48 hours (Steiner et al., 1998). Commonly isolated EA_ggEC serogroups include O44 and O3 (Steiner et al., 1998).

The enteropathogenic *E. coli* (EPEC) possess the *eae* gene, responsible for the attachment and effacement (A/E) lesions that occur on the intestinal lining of the host (Kaper et al., 2004). The EPEC group does not generally produce enterotoxins, but can

adhere to the intestinal mucosa (Levine et al., 1978). The EPEC are commonly observed in infant's diarrhea (Levine and Edelman, 1984). Common serogroups within the EPEC virulence group are O111 and O55 (Levine et al., 1978).

Enteroinvasive *E. coli* (EIEC) are extremely invasive serotypes of *E. coli* that utilizes adhesion proteins to bind and enter the intestinal cells (Resta-Lenert and Barrett, 2003). Although EIEC do not produce any toxins, the intestinal wall is severely damaged by the rapid colonization and implementation of adhesion proteins within the intestinal epithelial cells (Resta-Lenert and Barrett, 2001). Patients infected with EIEC usually have symptoms that include bloody voluminous diarrhea, blood and mucus in the stool, abdominal cramps, vomiting, fever, chills and malaise (Simonovic et al., 2000). Common serogroups of EIEC include O124, O28 and O112 (Pupo et al., 1997).

Enterohemorrhagic *E. coli* (EHEC) are the most widely known virulence group. This group of *E. coli* possesses the *eae* gene that produces the A/E lesions responsible for producing shiga-like toxins. The EHEC virulence group utilizes the protein, *intimin*. *Intimin* aids in the adhesion of A/E lesions to the microvillus of the intestinal wall by intimately attached to the translocated intimin receptor (*tir*). Susceptible cells within the large intestine have shiga toxin receptors that allow binding of the shiga toxins. These receptors are the globotriaosylceramide (Gb₃) receptors, part of the glycolipid globotriaosylceramide family. Monogastrics, including humans, have Gb₃ receptors whereas cattle lack the Gb₃ receptor (Pruimboom-Brees et al., 2000).

The Gb₃ receptors are established within the entire GI tract (duodenum, ileum, jejunum, colon and cecum) and within the kidney (Hoey et al., 2002). When shiga toxins bind to the Gb₃ receptor, the shiga toxin is then able to enter the cell and inhibit protein synthesis (Ling et al., 1998). The adherence of the shiga toxin to the intestinal mucosa

causes a rearrangement of actin in the host cell, causing significant deformation and death of the host cells (Strockbine et al., 1986). The deformed cells are sloughed off the intestinal lining into the intestinal tract which leads to accumulation of fluids subsequently causing diarrhea (Griffin and Tauxe, 1991). The damage to the intestinal lining of the host may lead to hemorrhaging, evident by bloody diarrhea (Riley et al., 1983). In severe cases, EHEC may enter the bloodstream causing septicemia, renal failure, hemolytic uremic syndrome (HUS) or thrombotic thrombocytopenic purpura (Griffin and Tauxe, 1991). Common serogroups of EHEC are O157, O26, O145, O103 and O111 (Fagen et al., 1999).

***Escherichia coli* O157:H7**

The EHEC serotype *Escherichia coli* O157:H7 was first associated with a major foodborne outbreak in 1982 and has since been commonly linked to foodborne cases associated with the consumption of contaminated, undercooked meat (Wetzel and LeJeune, 2006). In 2011, there was an estimated 1 million pounds of ground beef recalled due to possible O157:H7 contamination (USDA-FSIS, 2011). In 2010, Scharff reported O157:H7 cost United States residents approximately \$1 billion in premature deaths, lost wages and hospital bills, totaling approximately \$15,000 per case of O157:H7 illness (Scharff, 2010). Severity of illness can affect cost as well, ranging from \$26 for an individual who did not receive medical care to \$6.2 million for a death of a patient that suffered from HUS (Frenzen et al., 2005).

According to the Center for Disease Control (CDC), 265,000 shiga toxin-producing *Escherichia coli* infections were documented in 2010, of which O157:H7 comprises 33% of these infections (CDC, 2011). The O157:H7 serotype is a virulent

serotype with an infectious dose of ≤ 50 cells (Besser et al., 1999). Children and the elderly are the most susceptible to infections due to their weakened immature immunological state (Fox et al., 2009). It is because of the severity associated with consumption of contaminated food products that the food and meat industries have invested more than \$2 billion in intervention strategies within the processing plant (Kay, 2003; Koohmaraie et al., 2005).

Infection due to O157:H7 generally occurs through ingestion of contaminated products. Although the most common route of human O157:H7 infection is through the consumption of raw or undercooked meat, infections can result from other sources as well. For instance water runoff from ruminant environments (Armstrong et al., 1996) contaminated with feces used to irrigate crops (i.e. spinach and lettuce), the products can potentially become contaminated. Direct human interactions within various environments, such as cattle or other livestock facilities (Wetzel and LeJeune, 2006), petting zoos and swimming pools have also been suggested as potential environments where inoculation of O157:H7 can occur (Rangel et al., 2005).

Shiga toxin-producing *Escherichia coli* (STEC)

While most research of enterohemorrhagic *E. coli* (EHEC) has traditionally focused on O157:H7, there has been an increase in outbreaks of non-O157 strains of shiga toxin-producing *E. coli* (STEC). In fact, the CDC estimates that nearly 77% of all *E. coli* associated infections are due to non-O157 serogroups. The six additional STEC serogroups that the food and meat industry are concerned with include: O26, O45, O103, O111, O121 and O145 (Paddock et al., 2011). It has been traditionally believed that these non-O157 serogroups grow similarly as O157:H7 (Beutin et al., 1994). However, a

study recently conducted within our laboratory group suggests that the responses of these non-O157 serogroups are not universally the same as O157:H7.

The characterization of STEC involves each serogroup having capabilities to produce shiga toxins (Fratmico et al., 2004). Shiga toxins are compound toxins consisting of one A subunit and five B subunits which are linked by a disulfide bond (Paton and Paton, 1998). The A subunit inhibits protein synthesis by removing an adenine from the ribosome (O'Brien et al., 1987). The B subunits bind to the glycolipid receptors on the surface of the target cell and aids in endocytosis (O'Brien et al., 1987). Once the shiga toxin crosses the epithelial barrier within its host, it will then enter the bloodstream and target the Gb₃ receptors, thus inhibiting protein synthesis and causing lysis of the cells (Paton and Paton, 1998).

The STEC can produce shiga toxin I (*stx*₁), and/or, shiga toxin II (*stx*₂). Both of which have virtually the same biological activities (Strockbrine et al., 1986). The *stx*₁ is serologically identical to the *Shigella dysenteriae* toxin (O'Brien et al., 1987). The *stx*₂ can be serologically differentiated from *stx*₁ and *Shigella dysenteriae*, although all biological activities are similar (Strockbrine et al., 1986). Studies have suggested that *E.coli* that produce *stx*₂ are more virulent than *E. coli* that produce *stx*₁ (Dorn et al., 1989; Wadolkowski et al., 1991; Arthur et al., 2002). The increase in virulence and association of *stx*₂ to HUS is thought to be from the increased transcription rate *in vivo*, as opposed to *stx*₁ and the combination of both (Paton and Paton, 1998).

The O26 STEC harbors the *stx*₁ gene most predominately, but some strains of O26 are able harbor the *stx*₂ or the combinations of both genes (Arthur et al., 2002). The *stx*₁ gene can be harbored by O103, O45, O111, O121 and O145. Some STEC can harbor both shiga toxins such as O45, O111, O111 and O145. Although able to harbor the

combination of both shiga toxins, O45 and O111 cannot exclusively harbor *stx*₂ alone (Kaper et al., 1998)

Certain strains of STEC are capable of producing A/E lesions on enterocytes that rearranges the host actin, disassembling the microvillus and forming actin pedestals (Francis et al., 1986). All genes, including shiga toxin genes required for the A/E adherence and lesions are located on a single pathogenicity island referred to as the locus for enterocyte effacement (LEE; Donnenberg et al., 1997). The LEE is a cluster of genes that encode for a type III secretion system which aids in expressing the *eae* genes (O'Brien et al., 1987). The type III secretion system secretes translocated effector proteins such as EspA, EspB and EspD which are responsible for the initiation of signal transduction events (Paton and Paton, 1998). The *eae* gene is responsible for encoding the outer membrane protein, *intimin* and its receptor *tir*, which the binding of both proteins is responsible for the intimate attachment to the enterocyte (O'Brien et al., 1987; Kenny et al., 1997). *Enterohemolysin* is expressed through the *eae* gene and lyses erythrocytes to acquire iron to support growth of STEC (Paton and Paton, 1998). These virulence factors are responsible for A/E of the intestinal surface and can aid in virulence, but are not necessarily the cause of severe human illnesses (Donnenburg, 1993). Although these factors are characteristics of O157:H7, research suggests these factors are more common in pathogenic non-O157 strains, such as STEC (Beutin et al., 1994).

In 2010, the incidences of O157:H7 decreased to reach the national target of ≤ 1 case per 100,000 (CDC, 2011). Implementing pathogen prevention strategies in the abattoir and further processing plants such as hide and carcass washes, antimicrobial treatments, HACCP policies, cleanliness and good sanitation practices have lead to decreased O157:H7 populations (Elder, 2000). While the O157:H7 related outbreaks in

humans and animals have decreased, there has been an observed increase in non-O157 incidences. In response to this increase in prevalence, starting June of 2012 the USDA-FSIS will classify the STEC serotypes O26, O45, O103, O111, O121 and O145 as adulterants in non-intact raw beef and prohibit sale of any contaminated products (USDA-FSIS, 2012).

Role of enterohemorrhagic *Escherichia coli* in ruminant animals

Enterohemorrhagic *E. coli* are capable of colonizing within the GI tract of many ruminants (Callaway, 2002). Cattle are considered a major reservoir of O157:H7 (Low et al., 2005; Hussein et al., 2006; Ferens and Hovde, 2011). Cattle are asymptomatic carriers and play a role in the epidemiology of human O157:H7 infections (Griffin and Tauxe, 1991; Sargeant et al., 2006). One fecal prevalence study's results have suggested that approximately 30% of cattle are carriers of O157:H7 (Elder et al., 2000; Callaway et al., 2002). However, this can vary due to individual animal and season (Elder et al., 2000).

Studies have suggested that the large intestinal environment of cattle is selective for O157:H7 growth (Enss et al., 1996; Fox et al., 2009). This selective environment is related to GI tract mucosal components: fucose, galactose, mannose, *N*-acetylgalactosamine, *N*-acetylglucosamine, galacturonic acid, glucuronic acid and gluconic acid (Peekhaus and Conway, 1998; Fox et al., 2009). Mucus and its glycoprotein constituent, mucin are secreted within the large intestine and are responsible for the defense mechanisms to prevent colonization of pathogenic bacteria. Although a study conducted by Enss et al. (1996) indicated that another STEC serogroup O55 (Ludwig et al., 1996), more specifically the O55:B5 strain was able to modify mucin,

thus creating a more favorable condition for growth of bacteria by aiding in A/E mechanisms. These results suggest that other STEC such as O157:H7 may alter mucin to increase specific carbohydrate constituents, hence permitting O157:H7 to survive in the large intestine of cattle (Fox et al., 2009). Research has suggested that these mucosal components promote colonization of *E. coli* (K-12 and F-18) in the mouse model, but not specifically O157:H7 (Peekhaus and Conway, 1998; Chang et al., 2004). Although a more recent study conducted by Fox et al. (2009) suggested that more specifically gluconic acid may serve as a growth prominent for O157:H7 in the large intestine of beef cattle. These data further enhance the hypothesis that the extracellular components of the mucosal cells, such as mucin are able to serve as growth factors for O157:H7 (Miranda et al., 2004).

Escherichia coli O157:H7 has also been isolated from the oral cavity (Keen and Elder, 2002), digesta (Laven et al., 2003), ruminal contents (Elder et al., 2000), and feces (Elder et al., 2002; Keen and Elder, 2002). Some studies have suggested that the colon (Grauke et al., 2002) and the lymphoid tissue located at the recto-anal junction (Naylor et al., 2003) may be the primary colonization sites within the ruminants. Researchers have suggested that the recto-anal junction is very closely involved with O157:H7 colonization (Greenquist et al., 2005; Davis et al., 2006; Lim et al., 2007) due to O157:H7 specificity for follicle-associated epithelium within the recto-anal junction (Naylor et al., 2003).

Various serogroups of STEC (O7, O91, O146, O113, etc.) have been isolated from beef cattle (Beutin et al., 1997; Leomil et al., 2003), manure (Lahti et al., 2002) and raw beef (Kahn et al., 2002). Hussein and Bollinger (2005) indicated that the prevalence of non-O157 STEC were increased compared to O157:H7 in packing plants, supermarkets and fast food restaurants.

***Escherichia coli* O157:H7 shedding in cattle**

Shedding of O157:H7 in cattle appears to very sporadic (i.e. animal, season, time) and widespread (Meyer-Broseta et al., 2001). The amount and frequency of O157:H7 shedding varies greatly due to individual animal and season (LeJeune et al., 2004; Stanford et al., 2005). Although a number of various factors affect pathogen shedding, the only factor that has consistently been repeated is season (Edrington et al., 2006). In shedding prevalence studies, results suggest that typically, there is an increase in fecal shedding in spring, with potential peaking at 80% in feedlot cattle in the summer months; then falling drastically in the fall months with least amount of fecal shedding, 5-10% in the winter months (Elder et al., 2000; Barkocy-Gallagher et al., 2003; Edrington et al., 2006). Outbreaks of O157:H7 in humans seem to shadow this seasonal trend, primarily occurring in the summer months (Rangel et al., 2005). Cattle that are considered shedders of O157:H7 shed at a rate of $<10^2$ CFU/g of feces (Omisakin et al., 2003), whereas cattle that are considered super shedders can shed at least 10^4 CFU/g of feces (Chase-Topping et al., 2008).

Prevalence of *Escherichia coli* O157:H7 on cattle hides

Cattle can become infected and continually shed O157:H7 in feces up to 49 days (Wang et al., 1996). Therefore, O157:H7 can spread to other cattle through physical contact; such as animal-to-animal or animal-to-environment (Collis et al., 2004; Wetzel and LeJeune, 2006). These two routes are frequently the source of spread within the feedlot and transportation (Cuesta Alonso et al., 2007). Studies have indicated an increase in the prevalence of O157:H7 on cattle hides, specifically during transportation (Collis et al., 2004; Stanford et al., 2011).

Cattle within confined spaces with greater numbers of cattle tend to have more filth and fecal matter on hides (Arthur et al., 2009). Cattle hides at the time of harvest have been suggested to be one of the major sources of O157:H7 contamination on beef carcasses due to the increase fecal matter on the hides arriving at the abattoir (Nou et al., 2003; Bosilevac et al., 2005; Arthur et al., 2007). It has been reported that concentrations of *E. coli*, including O157:H7, can range from 10^2 to 10^7 CFU/g of feces at harvest (Jordan and McEwen, 1998). Cattle with potentially contaminated hides that arrive at the harvesting facilities increase the risk of microbial contamination of the meat (Bacon et al., 2000; Elder et al., 2000; Midgley and Desmarchelier, 2001). Even though carcasses are considered sterile after the hide has been removed, further contamination can occur through fabrication, viscera removal (Tutenel et al., 2003) and through contact with contaminated workers (Brackett, 1999).

***Escherichia coli* pre-harvest intervention strategies**

Controlling and/or eliminating fecal shedding in cattle has been a major goal in pre-harvest food safety. Intervention strategies have been installed within harvesting facilities, but still do not completely guarantee elimination of O157:H7, thus why researchers have put an emphasis on pre-harvest intervention strategies. Currently there are no pre-harvest interventions strategies that have been successful because strategies have not been repeatable and consistent or cost effective to the producer. According to Callaway (2010), pre-harvest intervention strategies can be divided into three basic categories: management practices and transport, cattle water and feed management and live animal treatments.

Management practices and transport

Fundamental animal husbandry practices have long been essential to maximize production efficiency and ensure animal health and welfare. Although O157:H7 is commonly isolated from a host, O157:H7 can survive within the environment as well. Studies have reported that O157:H7 can survive anywhere from 3 to 130 days within the soil (Maule, 2000).

Biosecurity factors also play a role in the O157:H7 spread. Cattle are not the only animals that able to harbor O157:H7 and all EHEC, pest such as rodents, birds, insects and flies can also harbor O157:H7 and can aid in the spread and epidemiology (Rice et al., 2003; Wetzel and LeJune, 2006; Ahmad et al., 2007). Allowing other domesticated livestock animals on the farm (i.e. goats and sheep), pets (i.e. dogs and cats) and unintentional wild animals and fowl (i.e. deer, ducks, geese, and pheasants) have been associated with increased O157:H7 shedding within cattle (Synge et al., 2003; Gunn et al., 2007; French et al., 2010).

Subsequently, O157:H7 is commonly indicated within a group of cattle as opposed to an entire farm, therefore it is suggested that cattle be placed within contemporary groups throughout the duration on the farm. Overcrowding of cattle within a group or on a livestock trailer increases the possibility for fecal O157:H7 transmission. The trailers used during transportation of potential carriers of O157:H7 are effortless pathways for the spread of pathogens (Mather et al., 2007). The trailers used to transport cattle from the feedlot to packer are frequently reported to test positive for O157:H7 but washing trailers did not decrease prevalence (Cuesta Alonso et al., 2007; Reicks et al., 2007).

Cattle water and feed management

Water sources shared by cattle are a common place where O157:H7 can be isolated. *Escherichia coli* O157:H7 can survive for up to six months within water sources (Hancock et al., 1998; LeJeune et al., 2001). Chlorination of the water supply within cattle water troughs has been tested as a possible solution to controlling the spread of O157:H7. Consequently, outside environmental factors such as sunlight and organic matter contribute to the decrease of effectiveness of chlorination (LeJune et al., 2004). Chlorination has led to cattle refusal to drink treated water because of organic matter accumulation and the distinct chlorine smell and taste. Electrolyzing water is another intervention strategy implemented in cattle drinking water. This involves the electrolysis of deionized water with free-chlorine concentrations to decrease bacterial populations (Shimamune et al., 1996). Electrolyzing water has been suggested to reduce O157:H7 concentrations as an in-plant hide cleaning strategy (Bosilevac et al., 2005), but not as a water source strategy.

Dietary manipulation has been a pre-harvest area researched in an attempt to reduce the O157:H7 burden before transportation and at the abattoir. Previous research by Wolin (1969) suggested that increased concentrations of volatile fatty acids (VFA) and a decreased pH could possibly inhibit *E.coli* growth. The toxicity of VFA on *E.coli* and other bacteria is due to the anion accumulation within the cell membrane when the pH is decreased (Russell, 1992). Whole cottonseed has been linked to reduce O157:H7 shedding because of the increase in short-chain VFA concentrations observed (Garber et al., 1995). Corn silage, addition of animal by-products (i.e. blood meal, meat meal, feather meal; Herriott et al., 1998) barley (Dargatz et al., 1997), distiller's grains (Jacob et al., 2008; Yang et al., 2010) and brewer's grain (Dewell et al., 2005) are feed

ingredients that have been linked to increased fecal shedding of O157:H7 in cattle. The fecal shedding is suggested to increase with use of these products because of the decreased effect on the short-chain VFA concentrations.

There has also been extensive research in the last decade on whether concentrate or forage diets are advantageous at reducing shedding of O157:H7. Concentrates are typically fed to feedlot cattle which are observed to have the greatest frequency of shedders (Tkalcic et al., 2000). Steam-flaked concentrates have indicated to increase O157:H7 shedding, while dry rolled concentrates have reduced shedding (Fox et al., 2007). Interestingly, greater concentrations of O157:H7 were observed within the feces of concentrate-fed animals compared to forage-fed animals (Tkalcic et al., 2000). Forage-fed animals were observed to shed O157:H7 for longer periods of time; 60 days for forage-fed as opposed to 16 days grain-fed (Van Baale et al., 2004). The O157:H7 proliferation due to potential variation in VFA concentrations suggests that cattle fed concentrates shed greater concentrations of O157:H7 (Tkalcic et al., 2000).

The amount and quality of forage also is important in dictating the amount and frequency of shedding. Poor quality forages (i.e. straw) exhibited a faster and greater death rate of O157:H7 compared to superior quality forages (i.e. grass silage; Franz et al., 2005). When cattle are suddenly changed from a full grain diet to a full forage diet, O157:H7 populations decreased (Diez-Gonzalez et al., 1998).

Feed additives have been researched to attempt to reduce the populations and patterns of O157:H7 shedding. Plant based products such as phenolics, essential oils and tannins have been evaluated for O157:H7 shedding reduction and/or elimination. Min et al. (2007) reported that secondary metabolites from natural phenolic compounds exhibit bactericidal activities in gram-negative bacteria. Research conducted with tannins has

reported to decrease populations of O157:H7 *in vitro*; although only concentrations of generic *E. coli* within the live animal were reduced (Min et al., 2007). The Federal Drug Administration has recognized essential oils as generally safe for human and animal consumptions (FDA, 2004). Essential oils contain microbial activity and are common components responsible for fragrance (Benchaar et al., 2008). Essential oils from plants are added to diets in attempt to reduce O157:H7 concentrations because of the toxicity exhibited on the cell wall of gram-negative bacteria such as O157:H7 (Dusan et al., 2006). Essential oils have reduced O157:H7 populations within cattle, while increasing fermentation (Benchaar et al., 2008).

Probiotics have previously been used by researchers to enhance production and health within the livestock (Tournut, 1989). Probiotics have also been proposed as a method to reduce pathogenic bacteria. By definition, probiotics are the use of live or dead cultures of microorganisms to alter the microbial environment of the GI tract. Probiotics within the GI tract, reduce pathogen concentrations by binding to the intestinal epithelium, subsequently preventing pathogens from binding (Collins et al., 1999). There are a few probiotic varieties that have been produced to aid in the reduction of shedding: direct-fed microbials (DFM), competitive exclusion (CE) and prebiotics.

Direct-fed microbials are fed to ruminants to improve ruminal fermentation. Direct-fed microbials are comprised of live or dead yeast, fungal or bacterial culture fermentation end-products (Callaway, 2009). Previous research suggests that *Streptococcus bovis*, *Lactobacillus gallinarum* (Ohya et al., 2001), *Lactobacillus acidilacti*, *Lactobacillus acidophilus* (Brashears and Galyean, 2002) and *Pediococcus* (Rodriguez-Palacios et al., 2009), have decreased O157:H7 concentrations in cattle.

Although research in direct-fed microbials is ongoing, the limiting factor is cost to the producer.

The concept of CE includes introducing non-pathogenic bacteria of the same species to the GI tract with intentions of reducing pathogenic bacteria colonization and shedding within the animal (Nurmi et al., 1992). The mechanisms of CE to reduce bacterial populations are to first create a hostile environment for other bacteria to survive within, eliminate bacterial receptor sites, produce and secrete antimicrobial metabolites and select and competitively deplete essential nutrients for other bacteria (Rolfe, 1991). The use of CE has reduced O157:H7 shedding for O111:NM and O26:H11 (Zhao et al., 2003).

Prebiotics are organic compounds that are indigestible by animals. Prebiotics are able to pass through the GI tract until being utilized as nutrient sources by the microbial populations within a specific environment in the animal. Prebiotics provide a competitive environment by competing with the pathogenic bacteria for nutrients and binding sites in the intestine (Zopf and Roth, 1996). Research has proposed that a synbiotic relationship (coupling CE and prebiotics) could reduce O157:H7 concentrations; by combining mechanisms of a competitive environment, elimination of receptors, and production of antimicrobial and depletion of nutrients available to pathogens (Bomba et al., 2002). To date most probiotic treatments have not been implemented due to the great cost associated and lack of producer return.

Colicins are a toxic exoproteins produced by *E. coli* strains that contain very specific cell wall receptors that apply an inhibitory effect on O157:H7, thus destroying or hindering growth by destroying the cell wall (Konisky, 1982; Smarda and Smajs, 1998). Callaway et al. (2004) indicated that colicins were effective in reducing O157:H7 *in*

vitro, while Cutler et al. (2007) indicated that colicins were effective in preventing other *E. coli* besides O157:H7 *in vivo*. Also, colicins have been supplemented to cattle diets, and research has reported a 2 log₁₀ CFU/g in reduction of O157:H7 populations (Nandiwada et al., 2004). Further research in this area is currently on-going but major barriers must be surmounted. Possible barriers include the ability of colicins to become resist to certain strains overtime (Alonso et al., 2000), immunity proteins produced by certain O157:H7 strains against colicins (Murinda et al., 1998), GI tract degradation and the cost associated (Lui et al., 2011).

Live animal *Escherichia coli* O157:H7 treatments

Aside from dietary manipulation, researchers have also investigated live animal pre-harvest treatments. Vaccinations, bacteriophages and sodium chlorate are the primary live animal treatments that are under investigation. The *E.coli* and *Salmonella* vaccinations utilized within swine and poultry have been effective in reducing pathogens (Johansen et al., 2000). Siderophore receptor and porin (SRP) protein vaccines are another vaccination being researched. *Escherichia coli* O157:H7 releases siderophore proteins within the intestinal environment to acquire iron. The SRP protein vaccination targets siderophores to obstruct the iron acquisition, thus leading to O157:H7 death. Recent research has indicated that the SRP injection has reduced O157:H7 concentrations by 98% in the feces (Thomson et al., 2009). The SRP protein vaccination is often not utilized because the cost is too great for producers to implement within production systems.

The bacterial extract vaccination is a three-dose regiment produced from O157:H7 extracts, particularly type III secreted proteins. The host protein type III

secretions release the *tir*, an adhesion protein that will integrate into the cell membrane. Once *tir* protein is established on the O157:H7 cell membrane it becomes a receptor for *intimin*, another adhesion protein necessary for the intimate colonization of O157:H7 (DeVinney et al., 1999). Research has reported that the *tir-intimin* binding is vital in O157:H7 A/E of the intestine (DeVinney et al., 1999; Cornick et al., 2002). The type III secretion vaccination introduces *intimin*_{O157} antibodies within the host and have decrease A/E lesions and colonization of the intestine of pigs (Dean-Nystrom et al., 2002). Research has indicated that this vaccine reduced O157:H7 shedding within cattle (Potter et al., 2004) and proposed to decrease shedding (McNeilly et al., 2010). A study conducted by Smith et al. (2009) suggested a 92% less likely probability for cattle within feedlot environments to be colonized in the terminal rectum mucosa when administered as a two dose regimen of a type III secretion vaccine. Research suggests that type III secreted protein vaccines and SRP vaccines have reduced fecal prevalence of O157:H7 within cattle that are naturally exposed to O157:H7 (Snedeker et al., 2011). Although, the results of vaccinations are promising and are approved for use in the United States (SRP and bacterial extract vaccines) and Canada (bacterial extract vaccines), producer use will remain uncertain because of cost and economic return and lack of study consistency.

Bacteriophages (phages) are viruses that can invade bacteria within any organism. The idea behind implementing phages in pre-harvest food safety involves phages that have the ability to specifically target host bacteria. When present, phages take over the target bacteria and replicate, continually releasing daughter cells to repeat the process (Callaway et al., 2008b). Exponential growth will continue as long as the target bacteria are present. One limiting factor to phage use is that phages can easily develop resistance.

Bacteriophages have been implemented to decrease O157:H7 populations in the ruminant GI tract (Bach et al., 2003; Callaway et al., 2008b; Rozema et al., 2009) as well as for hide and surfaces washes. Although phage treatments have been successful in reducing O157:H7 population *in vivo* (Bach et al., 2009) implementation on the farm has not been successful due to the cost relative to the producer (Callaway et al., 2008b).

Sodium chlorate has enzymatic properties that allow for beneficial pre-harvest application. Through the nitrate reductase enzyme, chlorate has the ability to reduce nitrate to nitrite. Nitrite reductase also reduces chlorate to chlorite within the cell cytoplasm (Stewart, 1988). An increase in chlorite concentrations within the cytoplasm destroys O157:H7 (Stewart, 1988). The addition of sodium chlorate to ruminal fluid indicated a reduction in O157:H7 concentrations *in vitro* (Anderson et al., 2000). Sodium chlorate supplementations to water supplies have decreased O157:H7 throughout the GI tract in cattle as well as sheep (Anderson et al., 2002; Callaway et al., 2002; Callaway et al., 2003).

Citrus by-products

The citrus production generates by-products for various food and household products, such as pulp and peel that have nutritional value and exhibit antimicrobial properties. Citrus by-products have been added to diets of dairy and beef cattle because of palatability, nutritional value and the relative decreased cost (Arthington et al., 2002). Citrus by-products are usually supplemented at a rate of 5-15% of the ration (Arthington et al., 2002). Citrus by-products are considered an energy, calcium, dietary fiber, and low protein concentrate ingredient, and are fed to cattle in the wet, dry, powdered or pelleted form (Arthington et al. 2002). Citrus by-products are readily available within the citrus-

producing regions such as California, Texas and Florida (Callaway et al., 2008a) and can be stored and transported long distances (Volanis et al., 2006).

Research has suggested that citrus by-products display antimicrobial activity against foodborne pathogens within the host animal (Dusan et al., 2006; Viuda-Martos et al., 2008; Friedly et al., 2009). Phytochemicals, such as limonene and citrullene are the most common and have the greatest microbial activity within citrus by-products (Di Pasqua et al., 2006). These oils are toxic to bacteria by increasing cell membrane permeability (Kim et al., 2005; Fisher and Phillips, 2006; Gill and Holley, 2006; Di Pasqua et al., 2007). The lipophilic properties of essential oils are able to permeabilize the cell wall and cytoplasmic membranes within gram-negative bacteria (Bakkali et al., 2006). The change in the fluidity of the membranes due to the permeabilization, allows essential oils to coagulate the cytoplasm (Gustafson et al., 1998), depleting ATP (Di Pasqua et al., 2006) and leading to lysis of the cell (Kim et al., 2005; Fisher and Phillips, 2006; Gill and Holley, 2006; Oussalah et al., 2006; Di Pasqua et al., 2007).

In vitro studies have suggested that the addition of >1% orange peel and pulp reduced O157:H7 and *Salmonella typhimurium* concentrations within rumen fluid (Callaway et al., 2008a). *In vivo* research conducted with sheep indicated that when feeding a 50/50 ration of orange peel and citrus pellets, the ration reduced inoculated O157:H7 and *S. typhimurium* populations in the rumen (Callaway et al., 2011a; Callaway et al., 2011b). Feeding orange peel and pulp to weanling pigs decreased ileal and cecal populations of diarrheagenic *E. coli*, while completely eliminating concentrations within the rectum (Collier et al., 2010). The *in vitro* and sheep *in vivo* data suggest that citrus by-products can reduce O157:H7 populations within cattle and sheep, more specifically within the rumen. Even though O157:H7 can reside in the rumen, the intestinal tract is

where colonization and shedding occurs (Laven et al., 2003). If citrus by-products can reduce O157:H7 populations within the rumen, it could subsequently reduce populations within the intestinal tract. This could make citrus by-products a dual purpose pre-harvest intervention strategy.


Future of pre-harvest

There have been numerous developments in pre-harvest intervention techniques. These strategies have been implemented on the farm and through the environment. Although cases of O157:H7 incidences have decreased, there has not been a strategy that has proven to completely eliminate O157:H7. The main limitation to pre-harvest food safety intervention techniques is the inability to successfully track pathogens. Biophotonics has been an area that researchers suggest could aid in the knowledge and pathogenesis of bacteria. Biophotonics introduces the idea of bacteria transformed with bioluminescent plasmids that can be detected under a specialized camera. Implementing the use of bioluminescence within the live animal model can serve as a research tool for comprehending the pathogenesis and physiological systems of healthy or diseased animals (Ryan et al., 2011). One of the main limitations to pre-harvest intervention strategies is the inability to recognize and diagnose an individual or a group of cattle that are carriers of O157:H7. Being able to physically track pathogens like O157:H7 will help researchers better understand what physiological events occur *in vivo*, allowing researchers to advance pre-harvest pathogen intervention strategies. Imaging of bioluminescent plasmids offers a rather straightforward, strong, lucrative and extremely sensitive resource for capturing biological *in vivo* processes (Dothager et al., 2009).

Biophotonics and bioluminescence

Biophotonics involves the introduction of a vector containing the *luxCDABE* operon, which encodes enzymes responsible for synthesis of the luciferase enzyme and the aldehyde substrate (Meighen, 1993) into the bacterial strain of interest. Biophotonics combines the scientific fields of biology, photonics and optical imaging.

Bioluminescence is the voluntary or involuntary emission of visible light in living organisms that is arbitrated to an enzyme catalyst (Meighen, 1993). An oxidized reaction that involves a luciferase enzyme and a substrate is produced and thus emits light (Zhao et al., 2004). Unique to bioluminescent bacteria is the ability to self-synthesize all substrates within the bacteria themselves for the production of light (Close et al., 2010). In order to generate light, the luciferase protein first binds to the two naturally occurring products within the cell; reduced riboflavin phosphate (FMNH₂) and oxygen (O₂). After the protein binds to each individually, it then binds to the synthesized long chain endogenous aliphatic aldehyde, produced by lipid biogenesis. This permits the *lux* cassette to implement the endogenous materials within the cell to form intermediates, in turn oxidizing the intermediates to produce emission of a blue-green light (Meighen, 1991). The overall reaction that is observed can be reviewed as:



The light that is emitted can be detected using an intensified charge-coupled device camera (Contag et al., 1995) and this camera can detect as few as 50 colonies (Siragusa et al., 1999).

Bacteria transformed with photonic plasmids that constitutively express of *luxCDABE* operon do not exhibit alterations in growth kinetics (Beyer and Bohm, 1996), biochemical mechanisms, serology, or structure in comparison to non-transformed strains

(Chen and Griffiths, 1996). Bioluminescence from bacterial luciferase also correlates well with *in vitro* enzyme activity, as the amount of luciferase protein and *lux* mRNA generated directly correlates with cell viability (Close et al., 2010; Siragusa et al., 1999) bacterial plate counts (Siragusa et al., 1999) and intracellular numbers of bacteria in cell culture (Maurer et al., 2000).

Successful models

Recent studies have adapted this bioluminescent technology for investigating a variety of physiologically relevant systems, including single, living cells (Willard et al., 1999), whole plants (Anderson et al., 1995), *Drosophila* (Brandes et al., 1996), and rodents (Contag et al., 1995, 1997). The ability to track bioluminescent pathogens *in vivo* through the GI tract has previously been successful in pigs, mice and fish (Willard et al., 2002; Burns-Guydish et al., 2005; Karsi et al., 2006; Moulton et al., 2009a). Successful *in vivo* tracking of *Salmonella typhimurium* transformed with the pAK1-*lux* bioluminescent plasmid was achieved within the swine (Willard et al., 2002; Moulton et al., 2009a) and *S. typhimurium* and *E. coli* transformed with various *luxCDABE* operons within the murine (Burns-Guydish et al., 2005; Foucault et al., 2010) GI tract models. *Ex vivo* tracking of *E. coli* transformed with Xen-14 plasmid has been successful in the bovine (Curbelo et al., 2010) and *E. coli*+pAK1-*lux* within the ovine (Moulton et al., 2009a), gerbils (Disson et al., 2008) and equine reproductive tracts (Ryan et al., 2010).

Advantages of bioluminescence

Real-time imaging using these luciferase reporter genes can aid in pre-harvest pathogen intervention *in vivo* by reducing the number of animals used in studies and providing answers to the pathogenesis of bacteria (Ryan et al., 2005). Luciferase reporter

genes applied to bioluminescent imaging are used to help monitor real-time physiological events related to colonization and pathogenesis of bacterial pathogens (Klerk et al., 2007; Luker and Luker, 2008; Ryan, 2011). Employing bioluminescence in live animal models will facilitate tracking of pathogens (Contag et al., 1997) and indicate the physical location of pathogens (Siragusa et al., 1999), which will increase researchers' understanding of pathogenic processes (Contag et al., 1995). Live animal models, which include *in vivo* cellular and molecular imaging, are being implemented through new technologies (Contag, 2002). Although technology like biophotonics does not seem to be appropriate for cattle producers or consumer use, it has the ability to help pre-harvest researchers in the pathogenic bacteria analysis and mode of action and transmission (Moulton et al., 2009a).

Literature cited

- Ahmad A, Nagaraja TG, Zurek L. Transmission of *Escherichia coli* O157:H7 to cattle by house flies. *Prev Vet Med* 2007; 80:74-81.
- Alonso G, Vilchez G, Vidal RL. How bacteria protect themselves against channel-forming colicins. *Inter Microbiol* 2000; 3:81-88.
- Armstrong GL, Hollingsworth J, Morris JG. Emerging foodborne pathogens: *Escherichia coli* O157:H7 as a model of entry of a pathogen into the food supply of the developed world. *Epidemiol* 1996; 18:24-51.
- Anderson RC, Buckley SA, Kubena LF, Stanker LH, Harvey RB, Nisbet DJ. Bactericidal effect of sodium chlorate on *Escherichia coli* O157:H7 and *Salmonella typhimurium* DT104 in rumen contents *in vitro*. *J Food Prot* 2000; 63:1038-1042.
- Anderson RC, Callaway TR, Anderson TJ, Kubena LF, Keith NK, Nisbet DJ. Bactericidal effect of sodium chlorate on *Escherichia coli* concentrations in bovine ruminal and fecal contents *in vivo*. *Microb Ecol Health Dis* 2002; 14:24-29.
- Anderson SL, SA Kay. Functional dissection of circadian clock- and phy- tochrome regulated transcription of the Arabidopsis CAB2 gene. *Proc Natl Acad Sci USA* 1995; 92:1500-1504.
- Angert E. Alternatives to binary fission in bacteria. *Nature Rev* 2005; 3:214-224.
- Arthington JD, Kunkle WE, Martin AM. Citrus pulp for cattle. 2002; p.317-328. In G. Rogers, and M. Poore (ed.), *The Veterinary Clinics of North America-Food Animal Practice*, WB Saunders Company, Philadelphia, PA.
- Arthur TM, Barkocy-Gallagher GA, Rivera-Betancourt M, Koohmaraie M. Prevalence and characterization of non-O157 shiga toxin-producing *Escherichia coli* on carcasses in commercial beef cattle processing plants. *Appl Environ Microbiol* 2002; 68:4847-4852.
- Arthur TM, Bosilevac JM, Brichta-Harhay DM, Guerini MN, Kalchayanand N, King DA, Shackelford SD, Wheeler TL, Koohmaraie M. Transportation and lairage environment effects on prevalence, numbers, and diversity of *Escherichia coli* O157:H7 on hides and carcasses of beef cattle at processing. *J Food Prot* 2007; 70:280-286.

- Arthur TM, Bosilevac JM, Brichta-Harhay DM, Kalchayanand N, Shackelford SD, Wheeler TL, Koohmaraie M. Source tracking of *Escherichia coli* O157:H7 and *Salmonella* contamination in the lairage environment at commercial U.S. beef processing plants and identification of an effective intervention. *J Food Prot* 2008; 71:1752-1760.
- Arthur TM, Keen JE, Bosilevac JM, Brichta-Harhay DM, Kalchayanand N, Shackelford SD, Wheeler TL, Nou X, Koohmaraie M. Longitudinal study of *Escherichia coli* O157:H7 in a beef cattle feedlot and role of high-level shedders in hide contamination. *Appl Environ Microbiol* 2009; 75:6515-6523.
- Bach SJ, McAllister YA, Veira DM, Gannon VPJ, Holley RA. Effect of bacteriophage DC22 on *Escherichia coli* O157:H7 in an artificial rumen system (RUSITEC) and inoculated sheep. *Amin Res* 2003; 52:89-101.
- Bach SJ, Johnson RP, Stanford K, McAllister TA. Bacteriophages reduce *Escherichia coli* O157:H7 levels in experimentally inoculated sheep. *Can J Anim Sci* 2009; 89:285-293.
- Bacon RT, Belk KE, Sofos JN, Clayton RP, Reagan JO, Smith GC. Microbial populations on animal hides and beef carcasses at different stages of slaughter in plants employing multiple-sequential interventions for decontamination. *J Food Prot* 2000; 63:1080-1086.
- Bakkali F, Averbeck S, Averbeck D, Idaomar M. Biological effects of essential oils-a review. *Food and Chem Tox* 2006; 46:446-475.
- Barkocy-Gallagher GA, Arthur TM, Rivera-Betancourt M, Nou X, Shackelford SD, Wheeler TL, Koohmaraie M. Seasonal prevalence of shiga toxin-producing *Escherichia coli*, including O157:H7 and non-O157 serotypes, and *Salmonella* in commercial beef processing plants. *J Food Prot* 2003; 66:1978-1986.
- Baranyi, J. Stochastic modelling of bacterial lag phase. *Int J Food Micro* 2002; 73:203-206.
- Benchaar C, Calsamiglia S, Chaves AV, Fraser GR, Colombatto D, McAllister TA, Beauchemin KA. A review of plant-derived essential oils in ruminant nutrition and production. *Anim Feed Sci Technol* 2008; 145:209-228.
- Besser RE, Griffin PM, Slutsker L. *Escherichia coli* O157:H7 gastroenteritis and the hemolytic uremic syndrome, an emerging infectious disease. *Ann Rev Med* 1999; 50:355-367.

- Beutin L, Akeksic S, Zimmerman S, Gleier K. Virulence factors and phenotypical traits of verotoxigenic strains of *Escherichia coli* isolated from human patients in Germany. *Med Microbiol Immunol (Berl)* 1994; 183:13-21.
- Beutin L, Geier D, Zimmermann S, Aleksic S, Gillespie HA, Whittam TS. Epidemiological relatedness and clonal types of natural population of *Escherichia coli* strains producing shiga toxins in separate populations of cattle and sheep. *Appl Environ Microbiol* 1997; 63:2175-2180.
- Beyer W, Böhm R. Labeling *Salmonella* live vaccine strains with the *lux* operon from *Vibrio fischeri* improves their detection and decimation from wild type. *Microbiol Res* 1996; 151:407-419.
- Bomba A, Nemcova R, Mudronova D, Guba P. The possibilities of potentiating the efficacy of probiotics. *Trends Food Sci Technol* 2002; 13:121-126.
- Bosilevac JM, Shackelford SD, Brichta DM, Koohmaraie M. Efficacy of ozonated and electrolyzed oxidative waters to decontaminate hides of cattle before slaughter. *J Food Prot* 2005; 68:1393-1398.
- Brackett RE. Incidence, contributing factors, and control of bacterial pathogens in produce. *Postharvest Bio Tech* 1999; 15:305-311.
- Brandes C, Plautz JD, Stanewsky R, Jamison CF, Straume M, Wood KV, Kay SA, Hall JC. Novel features of drosophila period transcription revealed by real-time luciferase reporting. *Neuron* 1996;16:687-692.
- Brashears MM, Galyean ML. Testing of probiotic bacteria for the elimination of *Escherichia coli* O157:H7 in cattle. Available at: <http://www.amif.org/PRProbiotics042302.htm>. Accessed Jan 2, 2012.
- Burns-Guydish SM, Olomu IN, Zhao H, Wong RJ, Stevenson DK, Contag CH. Monitoring age-related susceptibility of young mice to oral *Salmonella* enteric Serovar *typhimurium* infection using *in vivo* murine model. *Pediatr Res* 2005; 58:153-158.
- Callaway TR, Anderson RC, Genovese KJ, Poole TL, Anderson TJ, Byrd JA, Kubena LF, Nisbet DJ. Sodium chlorate supplementation reduces *E. coli* O157:H7 populations in cattle. *J Anim Sci* 2002; 80:1683-1689.
- Callaway TR, Edrington TS, Anderson RC, Genovese KJ, Poole TL, Elder RO, Byrd JA, Bischoff KM, Nisbet DJ. *Escherichia coli* O157:H7 populations in sheep can be reduced by chlorate supplementation. *J Food Prot* 2003; 66:194-199.

- Callaway TR, Stahl CH, Edrington TS, Genovese KJ, Lincoln LM, Anderson RC, Lonergan SM, Poole TL, Harvey RB, Nisbet DJ. Colicin concentrations inhibit growth of *Escherichia coli* O157:H7 *in vitro*. J Food Prot 2004; 67:2603-2607.
- Callaway TR, Edrington TS, Brabban AD, Keen JE, Anderson RC, Rossman ML, Engler MJ, Genovese KJ, Gwartney BL, Reagan JO, Poole TL, Harvey RB, Kutter EM, Nisbet DJ. Fecal prevalence of *Escherichia coli* O157, *Salmonella*, *Listeria*, and bacteriophage infecting *E. coli* O157:H7 in feedlot cattle in the southern plains region of the United States. Foodborne Pathog Dis 2006; 3:234-244.
- Callaway TR, Carroll JA, Arthington JD, Pratt C, Edrington TS, Anderson RC, Galyean ML, Ricke SC, Crandall P, Nisbet DJ. Citrus products decrease growth of *E. coli* O157:H7 and *Salmonella typhimurium* in pure culture and in fermentation with mixed ruminal microorganisms *in vitro*. Foodborne Path Dis 2008a; 5:621-627.
- Callaway TR, Edrington TS, Brabban AD, Anderson RC, Rossman ML, Engler MJ, Carr MA, Genovese KJ, Keen JE, Looper ML, Kutter EM, Nisbet DJ. Bacteriophage isolated from feedlot cattle can reduce *Escherichia coli* O157:H7 populations in ruminant gastrointestinal tracts. Foodborne Path Dis 2008b; 5:183-192.
- Callaway TR, Carr MA, Edrington TS, Anderson RC, Nisbet DJ. Diet, *Escherichia coli* O157:H7, and cattle: a review after 10 years. Curr Issues Mol Biol 2009; 11:67-80.
- Callaway TR. Pre-harvest control of *E. coli* O157:H7. In the National Cattlemen's Beef Association Beef Research and Knowledge Management Proceedings. 2010; Centennial CO.
http://www.beefresearch.org/CMDocs/BeefResearch/Safety/WhitePaper_proof8.pdf
- Callaway TR, Carroll JA, Arthington JD, Edrington TS, Rossman ML, Carr MA, Krueger MA, Ricke NA, Crandall SC, Nisbet DJ. *Escherichia coli* O157:H7 populations in ruminants can be reduced by orange peel product feeding. J Food Prot 2011a; 74:1917-1921.
- Callaway TR, Carroll JA, Arthington JD, Edrington TS, Rossman ML, Carr ML, Genovese KJ, Ricke SC, Crandall SC, Nisbet DJ. Orange peel pellets can reduce *Salmonella* populations in ruminants. Foodborne Path Dis 2011b; 8:1071-1075.
- CDC. 2011 Estimates of Foodborne Illnesses in the United States. Accessed December 23, 2011.
http://www.cdc.gov/nczved/divisions/dfbmd/diseases/ecoli_o157h7/#how_comm_on

- Chang DE, Smalley DJ, Tucker DL, Leatham MP, Norris WE, Stevenson SJ, Anderson AB, Grissom GE, Laux DC, Cohen PS, Conway T. Carbon nutrition of *Escherichia coli* in the mouse intestine. *Proc Natl Acad Sci USA* 2004; 101:7427-7432.
- Chase-Topping M, Gally D, Low C, Matthews L, Woolhouse M. Super-shedding and the link between human infection and livestock carriage of *Escherichia coli* O157. *Nat Rev Microbiol* 2008; 6:904-912.
- Chen J, Griffith MW. Salmonella Detection in Eggs Using *Lux+* Bacteriophages. *J Food Prot* 1996; 59:908-914.
- Close DM, Patterson SS, Ripp S, Baek SJ, Sanseverino J, Saylor GS. Autonomous bioluminescent expression of the bacterial luciferase gene cassette (*lux*) in a mammalian cell line. *PLoS ONE* 2010; 5: e12441.
- Collier CT, Carroll JA, Callaway TR, Arthington JD. Oral administration of citrus pulp reduces recovery of orally dosed *Escherichia coli* F18 in weaned pigs. *J Anim Vet Adv* 2010; 9:2140-2145.
- Collins DM, Gibson GR. Probiotics, prebiotics, and synbiotics: approaches for modulating the microbial ecology of the gut. *Amer J Clin Nutr* 1999; 69:1052S-1057S.
- Collis VJ, Reid CA, Hutchinson MJ, Davies MH, Wheeler PA, Small A, Buncic S. Spread of marker bacteria from the hides of cattle in a simulated livestock market and at an abattoir. *J Food Prot* 2004; 67:2397-2402.
- Contag CH, Contag PR, Mullins JI, Spillman SD, Stevenson DK, Benaron DA. Photonic detection of bacterial pathogens in living hosts. *Mol Microbiol* 1995; 18:593-603.
- Contag CH, Spillman SD, Contag PR, Oshiro M, Eames B, Dennery P, Stevenson DK, Benaron DA. Visualizing gene expression in living mammals using a bioluminescent reporter. *Photochem Pathobiol* 1997; 66:523-531.
- Contag PR. Whole-animal cellular and molecular imaging to accelerate drug development. *Drug Discov Today* 2002; 7:555-562.
- Cornick NA, Booher SL, Moon HW. Intimin facilitates colonization by *E. coli* O157:H7 in adult ruminants. *Infect Immun* 2002; 70:2704-2707.
- Cuesta Alonso EP, Gilliland SE, Krehbiel CR. Incidence and toxin production ability of *Escherichia coli* O157:H7 isolated from cattle trucks. *J Food Prot* 2007; 70:2383-2385.

- Curbelo J, Moulton K, Willard S. Photonic characteristics and *ex vivo* imaging of *Escherichia coli*-Xen14 within the bovine reproductive tract. *Therio* 2010; 73:48-55.
- Cutler SA, Lonergan SM, Cornick N, Johnson AK, Stahl CH. Dietary inclusion of colicin E1 is effective in preventing postweaning diarrhea by F18-positive *Escherichia coli* in pigs. *Antimicrob Ag Chemother* 2007; 51:3830-3835.
- Dargatz DA, Wells SJ, Thomas LA, Hancock DD, Garber LP. Factors associated with the presence of *Escherichia coli* O157 in feces of feedlot cattle. *J Food Prot* 1997; 60:466-470.
- Davis MA. Comparison of cultures from rectoanal-junction mucosal swabs and feces for detection of *Escherichia coli* O157 in dairy heifers. *Appl Environ Microbiol* 2006; 72:3766-3770.
- Dean-Nystrom EA, Gansheroff LJ, Mills M, Moon HW, O'Brien AD. Vaccination of pregnant dams with *intimin* (O157) protects suckling piglets from *E. coli* O157:H7 infection. *Infect Immun* 2002; 70:2414-2418.
- Delpy LP, Beranger G, Kaweh M. Bacterial growth phases. *Ann Inst Pasteur (Paris)* 1956; 91:112-115.
- DeVinney R, Stein M, Reinscheild D, Abe A, Ruschkowski S, Finlay BB. Enterohemorrhagic *E. coli* O157:H7 produces *Tir*, which is translocated to the host cell membrane but is not tyrosine phosphorylated. *Infect Immun* 1999; 7:2389-2398.
- Dewell GA, Ransom JR, Dewell RD, McCurdy K, Gardner IA, Hill AE, Sofos JN, Belk KE, Smith GC, Salman MD. Prevalence of and risk factors for *Escherichia coli* O157 in market-ready beef cattle from 12 US feedlots. *Foodborne Path Dis* 2005; 2:70-76.
- Diez-Gonzalez F, Callaway TR, Kizoulis MG, Russell JB. Grain feeding and the dissemination of acid-resistant *Escherichia coli* from cattle. *Science* 1998; 281:1666-1668.
- Di Pasqua R, Hoskins N, Betts G, Mauriello G. Changes in membrane fatty acids composition of microbial cells induced by addition of thymol, carvacrol, limonene, cinnamaldehyde, and eugenol in the growing media. *J Agric Food Chem* 2006; 54:2745-2749.
- Di Pasqua R, Betts G, Hoskins N, Edwards M, Ercolini D, Mauriello G. Membrane toxicity and antimicrobial compounds from essential oils. *J Agric Food Chem* 2007; 55:4863-4870.

- Disson O, Grayo S, Huillet E, Nikitas G, Langa-Vives F, Dussurget O, Ragon M, Le Monnier A, Babinet C, Cossart P, Lecuit M. Conjugated action of two species-specific invasion proteins for fetoplacental listeriosis. *Nature* 2008; 455:1114-1118.
- Donachie WD. Co-ordinate regulation of the *Escherichia coli* cell cycle or the cloud of unknowing. *Mol Microbiol* 2001; 40:779-785.
- Donnenberg MS, Tzipori S, McKee ML, O'Brien AD, Alroy J, Kaper JB. The role of the *eae* gene of enterohemorrhagic *Escherichia coli* in intimate attachment *in vitro* and in a porcine model. *J Clin Invest* 1993; 92:1418-1424.
- Donnenberg MS, Kaper JB, Finlay BB. Interactions between enteropathogenic *Escherichia coli* and host epithelial cells. *Trends Microbiol* 1997; 5:109-114.
- Dorn CR, Scotland SM, Smith HR, Willshaw GA, Rowe B. Properties of verocytotoxin-producing *Escherichia coli* of human and animal origin belonging to serotypes other than O157:H7. *Epidemiol Infect* 1989; 103:83-95.
- Dothager RS, Flentie K, Moss B, Pan MH, Kesarwala A, Piwnicka-Worms D. Advances in bioluminescent imaging of live animal models. *Curr Opin Biotech* 2009; 20:45-53.
- Doyle MP. *Escherichia coli* O157:H7 and its significance in foods. *Int J Food Microbiol* 1991; 12:289-302.
- Dusan F, Marian S, Katarina D, Dobroslava B. Essential oils-their antimicrobial activity against *Escherichia coli* and effect on intestinal cell viability. *Toxicol in Vitro* 2006; 20:1435-1445.
- Edrington TS, Callaway TR, Ives SE, Engler MJ, Loope ML, Anderson RC, Nisbet DJ. Seasonal shedding of *Escherichia coli* O157:H7 in ruminants: a new hypothesis. *Foodborne Pathog Dis* 2006; 3:413-421.
- Elder RO, Keen JE, Siragusa GR, Barkocy-Gallagher GA, Koohmaraire M, Laegreid WW. Correlation of enterohaemorrhagic *Escherichia coli* O157 prevalence in faeces, hides and carcasses of beef cattle during processing. *Proc Natl Acad Sci USA* 2000; 97:2999-3003.
- Elder RO, Keen JE, Wittum TE, Callaway TR, Edrington TS, Anderson RC, Nisbet DJ. Intervention to reduce fecal shedding of enterohemorrhagic *Escherichia coli* O157:H7 in naturally infected cattle using neomycin sulfate. *J Anim Sci* 2002; 80(Suppl 1):15(Abstr.).

- Enss ML, Muller H, Schmidt-Wittig U, Kownatzki R, Coenen M, Hedrich HJ. Effects of perorally applied endotoxin on colonic mucins of germfree rats. *Scand J Gastroenterol* 1996; 31:868-874.
- FDA, 2004. Food and Drug Administration of the US, 21 CFR 184. Accessed Jan 17, 2012. <http://www.cfsan.fda.gov/eafus.html>.
- Fisher K, Phillips CA. The effect of lemon, orange and bergamot essential oils and their components on the survival of *Campylobacter jejuni*, *Escherichia coli* O157, *Listeria monocytogenes*, *Bacillus cereus* and *Staphylococcus aureus* *in vitro* and in food systems. *J Appl Microbiol* 2006; 101:1232-1240.
- Foucault ML, Thomas L, Goussard S, Branchini BR, Grillot-Courvalin C. *In vivo* bioluminescence imaging for the study of intestinal colonization by *Escherichia coli* in mice. *Appl Environ Microbiol* 2010; 76:264-274.
- Fox JT, Depenbusch BE, Drouillard JS, Nagaraja TG. Dry-rolled or steam-flaked grain-based diets and fecal shedding of *Escherichia coli* O157 in feedlot cattle. *J Anim Sci* 2007; 85:1207-1212.
- Fox JT, Drouillard JS, Shi X, Nagaraja TG. Effects of mucin and its carbohydrate constituents on *Escherichia coli* O157 growth in batch culture fermentations with ruminal or fecal microbial inoculum. *J Anim Sci* 2009; 87:1304-1313.
- Francis DH, Collins JE, Duimstra JR. Infection of gnotobiotics pigs with an *Escherichia coli* O157:H7 strain associated with an outbreak of hemorrhagic colitis. *Infect Immun* 1986; 51:953-956.
- Franz E, van Diepeningen AD, de Vos OJ, van Bruggen AHC. Effects of cattle feeding regimen and soil management type on the fate of *Escherichia coli* O157:H7 and *Salmonella* enteric serovar *Typhimurium* in manure, manure-amended soil, and lettuce. *Appl Environ Microbiol* 2005; 71:6165-6174.
- French E, Rodriguez –Palacios A, LeJeune JT. Enteric bacterial pathogens with zoonotic potential isolated from farm-raised deer. *Foodborne Path Dis* 2010; 7:1031-1037.
- Frenzen PD, Drake A, Angulo FJ, The emerging infections program foodnet working group. Economic cost of illness due to *Escherichia coli* O157 infections in the United States. *J Food Prot* 2005; 68:2623-2630.
- Friedly EC, Crandall PG, Ricke SC, Roman M, O’Bryan CA, Chalova VI. *In vitro* anti-listerial effects of citrus oil fractions in combination with organic acids. *J Food Sci* 2009; 74:M67-M72.

- Garber LP, Wells SJ, Hancock DD, Doyle MP, Tuttle J, Shere JA, Zhao T. Risk factors for fecal shedding of *Escherichia coli* O157:H7 in dairy calves. *J Amer Vet Med Assoc* 1995; 207:46-49.
- Gill AO, Holley RA. Disruption of *Escherichia coli*, *Listeria monocytogenes*, and *Lactobacillus sakei* cellular membranes by plant oil aromatics. *Int J Food Microbiol* 2006; 108:1-9.
- Griffin PM, Tauxe RV. The epidemiology of infections caused by *Escherichia coli* O157:H7, other enterohemorrhagic *E. coli* and the associated hemolytic uremic syndrome. *Epidemiol Rev* 1991; 13:60-98.
- Grauke LJ, Kudva IT, Yoon JW, Hunt CW, Williams CJ, Hovde CJ. Gastrointestinal tract location of *Escherichia coli* O157:H7 in ruminants. *Appl Environ Microbiol* 2002; 68:2269-2277.
- Greenquist MA, Drouillard JS, Sargeant JM, Depenbusch BE, Shi X, Lechtenberg KF, Nagaraja TG. Comparison of rectoanal mucosal swab cultures and fecal cultures for determining prevalence of *Escherichia coli* O157:H7 in feedlot cattle. *Appl Environ Microbiol* 2005; 71:6431-6433.
- Guarner F, Malagelada JR. Gut flora in health and disease. *The Lancet* 2003; 361:512-519.
- Gunn GJ, McKendrick IJ, Ternent HE, Thomson-Carter F, Foster G, Synge BA. An investigation of factors associated with the prevalence of verocytotoxin producing *Escherichia coli* O157 shedding in Scottish beef cattle. *Vet J* 2007; 174:554-564.
- Gustafson JE, Liew YC, Chew S, Markham JK, Bell HC, Wyllie SG, Warmington JR. Effects of tea oil on *Escherichia coli*. *Lett Appl Microbiol* 1998; 26:194-198.
- Herriott DE, Hancock DD, Ebel ED, Carpenter LV, Rice DH, Besser TE. Association of herd management factors with colonization of dairy cattle by shiga toxin-positive *Escherichia coli* O157. *J Food Prot* 1998; 61:802-807.
- Hilborn, ED, Mermin JH, Mshar PA, Hadler JL, Voetsch A, Wojtkunski CW, Swartz M, Mshar R, Lambert-Fair MA, Farrar JA, Glynn MK, Slutsker L. A multistate outbreak of *Escherichia coli* O157:H7 infections associated with consumption of mesclun lettuce. *Arch Intern Med* 1999; 159:1758-1764.
- Hoey ED, Currie C, Else RW, Nutikka A, Lingwood CA, Gally DL, Smith DGE. Expression of receptor for verotoxins 1 from *Escherichia coli* O157 on bovine intestinal epithelium. *J Med Microbiol* 2002; 51:143-149.

- Hudault S, Guignot J, Servin AL. *Escherichia coli* strains colonizing the gastrointestinal tract protect germfree mice against *Salmonella typhimurium* infection. *Gut* 2000; 49:47-55.
- Hussein HS, Sakuma T. Prevalence of shiga toxin-producing *Escherichia coli* in dairy cattle and their products. *J Dairy Sci* 2005; 88:450-465.
- Hussein HS. Prevalence and pathogenicity of shiga toxin-producing *Escherichia coli* in beef cattle and their products. *J Anim Sci* 2006; 85:E63-E72.
- Jacob ME, Fox JT, Drouillard JS, Renter DG, Nagaraja TG. Effects of dried distillers' grain on fecal prevalence and growth of *Escherichia coli* O157 in batch culture fermentations from cattle. *Appl Environ Microbiol* 2008; 74:38-43.
- Jensen PR, Hammer K. Minimal requirements for exponential growth of *Lactococcus lactis*. *Appl Environ Microbiol* 1993; 59:4363-4366.
- Johansen M, Andersen LO, Thomsen LK, Busch ME, Wachmann H, Jorsal SE, Gyles CL. Prevention of edema disease in pigs by passive immunization. *Can J Vet Res* 2000; 64:9-14.
- Johnson, JR. Virulence factors in *Escherichia coli* urinary tract infection. *Clin Microbiol Rev* 1991; 4:80-128.
- Jordan D, McEwen SA. Effect of duration of fasting and a short-term high-roughage ration on the concentration of *Escherichia coli* biotype 1 in cattle feces. *J Food Prot* 1998; 61:531-534.
- Kaper JB, O'Brien AD. *Escherichia coli* O157:H7 and Other Shiga Toxin-Producing *E. coli* Strains. Washington, DC: ASM Press; 1998.
- Kaper JB, Nataro JP, Mobley HLT. Pathogenic *Escherichia coli*. *Nature Rev Microbiol* 2004; 2:123-140.
- Karsi A, Menanteau-Ledouble S, Lawrence ML. Development of luminescent *Edwardsiella ictaluri* for non-invasive disease monitoring. *FEMS Microbiol Lett* 2006; 260:216-223.
- Kay S. *E. coli* O157:H7: The cost during the last 10 years. *Meat and Poultry News* 2003; 3:26-34.
- Keen JE, Elder RO. Isolation of shiga-toxigenic *Escherichia coli* O157 from hide surfaces and the oral cavity of finished beef feedlot cattle. *J Am Vet Med Assoc* 2002; 220:756-763.

- Kenny B, DeVinney R, Stein M, Reinscheild DJ, Frey EA, Finlay BB. Enteropathogenic *E.coli* (EPEC) transfers its receptor for intimate adherence into mammalian cells. *Cell* 1997; 91:511-520.
- Kim J, Marshall MR, Wei CI. Antimicrobial activity of some essential oil components against five foodborne pathogens. *J Agric Food Chem* 1995; 43:2839-2845.
- Klerk CPW, Overmeer RM, Niers TMH, Versteeg HH, Richel DJ, Buckle T, Van Noorden CJF, Van Tellingen O. Validity of bioluminescence measurements for noninvasive *in vivo* imaging of tumor load in small animals. *Biotechniques* 2007; 43(Supple.):7-13.
- Kolter R, Siegele DA, Tormo A. That stationary phase of the bacterial life cycle. *Ann Rev Microbiol* 1993; 47: 855-874.
- Konisky J. Colicins and other bacteriocins with established modes of action. *Ann Rev Microbiol* 1982; 36:125-144.
- Koohmaraie M, Arthur TM, Bosilevac JM, Guerini M, Shackelford SD, Wheeler TL. Post-harvest interventions to reduce/eliminate pathogens in beef. *Meat Sci* 2005; 71:79-91.
- Lahti E, Eklund M, Ruutu P, Siitonen A, Rantala L, Nuorti P, Honkanen-Buzalski T. Use of phenotyping and genotyping to verify transmission of *Escherichia coli* O157:H7 from dairy farms. *Eur J Clin Microbiol Infect Dis* 2002; 21:189-195.
- Laven RA, Ashmore A, Stewart CS. *Escherichia coli* in the rumen and colon of slaughter cattle, with particular reference to *E. coli* O157. *Vet J* 2003; 165:78-83.
- LeJeune JT, Besser TE, Hancock DD. Cattle water troughs as reservoirs of *Escherichia coli* O157. *Appl Environ Microbiol* 2001; 67:3053-3057.
- LeJeune JT, Besser TE, Rice DH, Berg JL, Stilborn RP, Hancock DD. Longitudinal study of fecal shedding of *Escherichia coli* O157:H7 in feedlot cattle: predominance and persistence of specific clonal types despite massive cattle population turnover. *Appl Environ Microbiol* 2004; 70:377-384.
- Leomil L, Aidar-Ugrinovich L, Guth BEC, Irino K, Vettorati MP, Onuma DL, de Castro AFP. Frequency of shiga toxin-producing *Escherichia coli* (STEC) isolates among diarrheic and non-diarrheic calves in Brazil. *Vet Microbiol* 2003; 97:103-109.

- Levine MM, Nalin DR, Hornick RB, Bergquist EJ, Waterman DH, Young CR, Sotman S, Rowe B. *Escherichia coli* strains that cause diarrhea but do not produce heat-labile or heat-stable enterotoxins and are non-invasive. *The Lancet* 1978; 1:1119-1122.
- Levine MM, Edelman R. Enteropathogenic *Escherichia coli* of classic serotypes associated with infant diarrhea: epidemiology and pathogenesis. *Epidemiol Rev* 1984; 6:31-51.
- Levine, MM. *Escherichia coli* that causes diarrhea: enterotoxigenic, enteropathogenic, enteroinvasive, enterohemorrhagic, and enteroadherent. *J Infect Dis* 1987; 155:377-389.
- Lim JY, Li J, Sheng H, Besser TE, Potter K, Hovde CJ. *Escherichia coli* O157:H7 colonization at the rectoanal junction of long-duration culture-positive cattle. *Appl Environ Microbiol* 2007; 73:1380-1382.
- Ling H, Boodhoo A, Hazes B, Cummings MD, Armstrong GD, Brunton JL, Read RJ. Structure of the shiga-like toxin I B-Pentamer complex with an analogue of its receptor Gb₃. *Biochemistry* 1998; 37:1777-1788.
- Low JC, McKendrick IJ, McKechnie C, Fenlon D, Naylor SW, Currie C, Smith DGE, Allison L, Gally DL. Rectal carriage of enterohemorrhagic *Escherichia coli* O157 in slaughtered cattle. *Appl Environ Microbiol* 2005; 71:93-97.
- Ludwig K, Bitzan M, Zimmerman S, Klothe M, Ruder H, Muller-Wiefel DE. Immune response to non-O157 verotoxin-producing *Escherichia coli* in patients with hemolytic uremic syndrome. *J Infect Dis* 1996; 174:1028-1039.
- Luker KE, Luker GD. Applications of bioluminescence imaging to antiviral research and therapy: Multiple luciferase enzymes and quantization. *Antiviral Res* 2008; 78:179-187.
- Maurer HH, Bickeboeller-Friedrich J, Kraemer T, Peters FT. Toxicokinetics and analytical toxicology of amphetamine-derived designed drugs ('Ecstasy'). *Toxicology Letters* 2000; 112-13:133-142.
- Mather AE, Innocent GT, McEwen SA, Reilly WJ, Taylor DJ, Steele WB, Gunn GJ, Ternent HE, Reid SWJ, Mellor DJ. Risk factors for hide contamination of Scottish cattle at slaughter with *Escherichia coli* O157. *Prev Vet Med* 2007; 80:257-270.
- Matic I, Radman M, Taddei F, Picard B, Doit C, Bingen E, Denamur E, Elion J. Highly variable mutation rates in commensal and pathogenic *Escherichia coli*. *Science* 1997; 277:1833-1834.

- Maule, A. Survival of verocytotoxigenic *Escherichia coli* O157 in soil, water and on surfaces. *J Appl Microbiol* 2000; 29:71S-78S.
- McNeilly TN, Mitchell MC, Rosser T, McAteer S, Low JC, Smith DGE, Huntley JF, Mahajan A, Gally DL. Immunization of cattle with a combination of purified *intimin-531*, *EspA* and *Tir* significantly reduces shedding of *Escherichia coli* O157:H7 following oral challenge. *Vaccine* 2010; 28:1422-1428.
- Meighen EA. Molecular biology of bacterial bioluminescence. *Microbiol Rev* 1991; 55:123-142.
- Meighen EA. Bacterial bioluminescence: organization, regulation, and application of the *lux* genes. *The FASEB Journal* 1993; 7:1016-1022.
- Meyer-Broseta S, Bastian SN, Arne PD, Cerf O, Sanaa M. Review of epidemiological surveys on the prevalence of contamination of healthy cattle with *Escherichia coli* serogroup O157:H7. *Int J Hgy Environ Health* 2001; 203:347-361.
- Midgley J, Desmarchelier P. Pre-slaughter handling of cattle and shiga toxin-producing *Escherichia coli* (STEC). *Lett Appl Microbiol* 2001; 32:307-311.
- Min BR, Pinchak WE, Anderson RC, Callaway TR. Effect of tannins on the *in vitro* growth of *Escherichia coli* O157:H7 and *in vivo* growth of generic *Escherichia coli* excreted from steers. *J Food Prot* 2007; 70:543-550.
- Miranda RL, Conway T, Leatham MP, Chang DE, Norris WE, Allen JH, Stevenson SJ, Laux DC, Cohen PS. Glycolytic and gluconeogenic growth of *Escherichia coli* O157:H7 (EDL933) in the mouse intestine. *Infect Immun* 2004; 73:1666-1676.
- Moseley SL, Huq I, Alim ARMA, So M, Samadpour-Motalebi M, Falkow S. Detection of enterotoxigenic *Escherichia coli* by DNA colony hybridization. *J Infect Dis* 1980; 142:892-898.
- Moulton K, Ryan P, Lay D, Willard S. Postmortem photonic imaging of *lux*-modified *Salmonella typhimurium* within the gastrointestinal tract of swine following oral inoculation *in vivo*. *J Anim Sci* 2009a; 87: 2239-2244.
- Moulton K, Ryan P, Christiansen D, Hopper R, Klausner C, Bennett W, Rodta-Palenik, Willard ST. *Ex Vivo* bioluminescence imaging of late gestation ewes following intrauterine inoculation with *lux*-modified *Escherichia coli*. *Comp Immunol Microbiol Infect Dis* 2009b; 32:429-438.
- Murinda SE, Liu SM, Roberts RF, Wilson RA. Colicinogeny among *Escherichia coli* serotypes, including O157:H7 representing four closely related diarrheagenic clones. *J Food Prot* 1998; 61:1431-1438.

- Nagy B, Fekete PZ. Enterotoxigenic *E.coli* (ETEC) in farm animals. *Vet Res* 1999; 30:259-284.
- Nandiwada LS, Schamberger GP, Schafer HW, Diez-Gonzales F. Characterization of an E2-type colicin and its application to treat alfalfa seeds to reduce *Escherichia coli* O157:H7. *Int J Food Microbiol* 2004; 93:267-279.
- Nataro J, Yikang D, Cookson S, et al. Heterogeneity of enteroaggregative *Escherichia coli* virulence demonstrated in volunteers. *J Infect Dis* 1995; 171:465-468.
- Naylor SW, Low JC, Besser TE, Mahajan A, Gunn GJ, Pearce MC, McKendrick IJ, Smith DGE, Gally DL. Lymphoid follicle-dense mucosa at the terminal rectum is the principal site of colonization of enterohemorrhagic *Escherichia coli* O157:H7 in the bovine host. *Infect Immun* 2003; 71:1505-1512.
- Nou X, Rivera-Betancourt M, Bosilevac JM, Wheeler TL, Shackelford SD, Gwarty BL, Reagan JO, Koohmaraie M. Effect of chemical dehairing on the prevalence of *Escherichia coli* O157:H7 and the levels of aerobic bacteria and *Enterobacteriaceae* on carcasses in a commercial beef processing plant. *J Food Prot* 2003; 66:2005-2009.
- Nurmi E, Nuotio L, Schncitz C. The competitive exclusion concept: development and future. *Int J Food Microbiol* 1992; 15:237-240.
- O'Brien AD, Holmes RK. Shiga and shiga-like toxins. *Microbiol Rev* 1987; 51:206-220.
- Ohya T, Akiba M, Ito H. Use of a trial probiotic product in calves experimentally infected with *Escherichia coli* O157. *Japan Agric Res Quart* 2001; 35:189-194.
- Omisakin F, MacRae M, Ogden ID, Strachan NJC. Concentration and prevalence of *Escherichia coli* O157 in cattle feces at slaughter. *Appl Environ Microbiol* 2003; 69:2444-2447.
- Oussalah M, Caillet S, Lacroix M. Mechanisms of action of Spanish oregano, Chinese cinnamon, and savory essential oils against cell membranes and walls of *Escherichia coli* O157:H7 and *Listeria monocytogenes*. *J Food Prot* 2006; 69:1046-1055.
- Paddock Z, Shi X, Bai J, Nagaraja TG. Applicability of a multiplex PCR to detect O26, O45, O111, O121, O145, and O157 serogroups of *Escherichia coli* in cattle feces. *Vet Microbiol* 2011.
- Paton JC, Paton AW. Pathogenesis and diagnosis of shiga toxin-producing *Escherichia coli* infections. *Clin Microbiol Rev* 1998; 11:450-479.

- Peekhaus N, Conway T. What's for dinner? Entner-Doudoroff metabolism in *Escherichia coli*. J Bacteriol 1998; 180:3495-3502
- Potter AA, Klashinsky S, Li Y, Frey E, Townsend H, Rogan D, Erickson GE, Hinkley S, Klopfenstein TJ, Moxley RA, Smith DT, Finlay BB. Decreased shedding of *Escherichia coli* O157:H7 by cattle following vaccination with type III secreted proteins. Vaccine 2004; 22:362-369.
- Pruimboom-Brees IM, Morgan TW, Ackermann MR, Nystrom ED, Samuel JE, Cornick NA, Moon HW. Cattle lack vascular receptors for *Escherichia coli* O157:H7 shiga toxins. PNAS 2000; 97:10325-10329.
- Pupo GM, Karaolis DK, Lan R, Reeves PR. Evolutionary relationships among pathogenic and nonpathogenic *Escherichia coli* strains inferred from multilocus enzyme electrophoresis and mdh sequence studies. Infect Immun 1997; 65:2685-2692.
- Rangel JM, Sparling PH, Crowe C, Griffin PM, Swerdlow DL. Epidemiology of *Escherichia coli* O157:H7 outbreaks, United States, 1982-2002. Emerg Infect Dis 2005; 11:603-609.
- Ray DE, Schaffer HD. E. coli again: A troubling new twist with serious consequences. Agricultural Policy Analysis Center, University of Tennessee, Knoxville, TN. Accessed 2 Feb 2012. <http://www.agpolicy.org/weekcol/567.html>
- Reicks AL, Brashears MM, Adams KD, Brooks JC, Blanton JR, Miller MF. Impact of transportation of feedlot cattle to the harvest facility on the prevalence of *Escherichia coli* O157:H7, *Salmonella*, and total aerobic microorganisms on hides. J Food Prot 2007; 70:17-21.
- Reid G, Howard J, Gan BS. Can bacterial interference prevent infection? Trends in Microbiology 2001; 9:424-428.
- Rembacken BJ, Snelling AM, Hawkey PM, Chalmers DM, Axon ATR. Non-pathogenic *Escherichia coli* versus mesalazine for the treatment of ulcerative colitis: a randomized trial. The Lancet 1999; 354:635-639.
- Resta-Lenert S, Barrett KE. Modulation of tight junction structure by enteroinvasive *E.coli* (EIEC) is prevented by H₂O₂ scavengers and protein kinase C (PKC) inhibitors. Gastroenterology 2001;120:3802.
- Resta-Lenert S, Barrett KE. Live probiotic protect intestinal epithelial cells from the effects of infection with enteroinvasive *Escherichia coli* (EIEC). Gut 2003; 52:988-997.

- Rice DH, Hancock DD, Besser TE. Fecal culture of wild animals for *Escherichia coli* O157:H7. *Vet Rec* 2003; 152:82-83.
- Riley LW, Remis RS, Helgerson SD, McGee HB, Wells JG, Davis BR, Hebert RJ, Olcott ES, Johnson LM, Hargrett NT, Blake PA, Cohen ML. Hemorrhagic colitis associated with a rare *Escherichia coli* serotypes. *N Engl J Med* 1983; 308:681-685.
- Rodriguez-Palacios A, Staempfli HR, Duffield T, Weese JG. Isolation of bovine intestinal *Lactobacillus plantarum* and *Pediococcus acidilactici* with inhibitory activities against *Escherichia coli* O157:H7 in feedlot cattle. *J Appl Microbiol* 2009; 106:393-401.
- Rolfe RD. Population dynamics of the intestinal tract. In: Colonization control of human bacterial enteropathogens in poultry. LC Blakenship (eds.) Academic Press, Inc., San Diego, CA. 1991: 59-75.
- Rolfe RD. The role of probiotic cultures in the control of gastrointestinal health. *J Nutr* 2000; 130(2S Suppl):396S-402S.
- Roszak DB, Colwell RR. Survival strategies of bacteria in the natural environment. *Microbiol Rev* 1987, 51:365-379.
- Rozema EA, Stephens TP, Bach SJ, Okine EK, Johnson RP, Stanford IM, McAllister TA. Oral and rectal administration of bacteriophages for control of *Escherichia coli* O157:H7 in feedlot cattle. *J Food Prot* 2009; 72:241-250.
- Russell JB. Another explanation for the toxicity of fermentation acids at low pH: anion accumulation versus uncoupling. *J Appl Bacterio* 1992; 73:363-370.
- Ryan PL, Youngblood RC, Harvill J, Willard ST. Photonic monitoring in real time of vascular endothelial growth factor receptor 2 gene expression under relaxin-induced conditions in a novel murine wound model. *Ann NY Acad Sci* 2005; 1041:398-414.
- Ryan PL, Christiansen DL, Hopper RM, Walters FK, Cooley AJ, Moulton K, Willard ST. Use of bioluminescence imaging technology and *lux*-modified bacteria to determine pathogen progression during uterine infections in the pregnant mare and rate of clearance post partum. *Anim Reprod Sci* 2010; 121S:S341-S342.
- Ryan PL, Christiansen DL, Hopper RM, Walters FK, Moulton K, Curbelo J, Greene JM, Willard ST. Horse Species Symposium: A novel approach to monitoring pathogen progression during uterine and placental infection in the mare using bioluminescence imaging technology and *lux*-modified bacteria. *J Anim Sci* 2011; 89:1541-1551.

- Sargeant JM, Amezcua MR, Rajic A, Waddell L. Pre-harvest interventions to reduce the shedding of *E. coli* O157 in the faeces of weaned domestic ruminants: a systematic review. *Zoo and Pub Health* 2006; 54:260-277.
- Scharff, RL 2010. Health-related costs from foodborne illness in the United States. 3 May, Accessed 2010.
<http://www.producesafetyproject.org/admin/assets/files/Health-Related-Foodborne-Illness-Costs-Report.pdf-1.pdf>.
- Sheng H, Lim JY, Watkins MK, Minnich SA, Hovde CJ. Characterization of an *Escherichia coli* O157:H7 O-antigen deletion mutant and effect of the deletion on bacterial persistence in the mouse intestine and colonization at the bovine terminal rectal mucosa. *Appl Environ Microbiol* 2008; 74:5015-5022.
- Shimamune T, Nakamatsu K, Sawamoto K, Nishiki K. Method for electrolyzing water. United States Patent #5,589,052. 1996.
- Simonovic I, Rosenberg J, Koutsouris A, et al. Enteropathogenic *Escherichia coli* dephosphorylates and dissociates occludin from intestinal epithelial tight junctions. *Cell Microbiol* 2000;2:305–15.
- Siragusa GR, Nawotka K, Spilman SD, Contag PR, Contag CH. Real-time monitoring of *Escherichia coli* O157:H7 adherence to beef carcass surface tissues with a bioluminescent reporter. *Appl Environ Microbiol* 1999; 65:1738-1745.
- Smarda J, Smajs D. Colicins-exocellular lethal proteins of *Escherichia coli*. *Folia Microbiol* 1998; 43:563-582.
- Smith DR, Moxley RA, Peterson RE, Klopfenstein TJ, Erickson GE, Bretschneider G, Berberov EM, Clowser S. A two-dose regimen of a vaccine against type III secreted protein reduced *Escherichia coli* O157:H7 colonization of the terminal rectum in beef cattle in commercial feedlots. *Foodborne Path Dis* 2009; 6:155-161.
- Snekeder KG, Campbell M, Sargeant JM. A systematic review of vaccination to reduce the shedding of *Escherichia coli* O157 in faeces of domestic ruminants. *Zoo Pub Health* 2011; 59:126-138.
- Stanford K, Bach SJ, Marx TH, Jones S, Hansen JR, Wallins GL, Zahiroddini H, McAllister TA. Monitoring *Escherichia coli* O157:H7 in inoculated and naturally colonized feedlot cattle and their environment. *J Food Prot* 2005; 68:26-33.
- Stanford K, Stephens TP, McAllister TA. Use of model super-shedders to define the role of pen floor and hide contamination in the transmission of *Escherichia coli* O157:H7. *J Anim Sci* 2011; 89:237-244.

- Steiner TS, Lima AAM, Nataro JP, Guerrant RL. Enteroaggregative *Escherichia coli* produce intestinal inflammation and growth impairment and cause interleukin-8 release from intestinal epithelial cells. *J Infect Dis* 1998; 177:88-96.
- Stewart V. Nitrate respiration in relation to facultative metabolism in enterobacteria. *Microbiol Rev* 1988; 52:190-232.
- Strockbine NA, Marques RM, Newland JW, Smith HW, Holmes RK, O'Brien AD. Two toxin-converting phages from *Escherichia coli* O157:H7 strain 933 encode antigenetically distinct toxins with similar biological activities. *Infect Immun* 1986; 53:135-140.
- Syngé BA, Chase-Toppings ME, Hopkins GF, McKendrick IJ, Thomson-Carter F, Gray Rusbridge SM, Munro FI, Foster G, Gunn GJ. Factors influencing the shedding of verocytotoxin-producing *Escherichia coli* O157 by beef suckler cows. *Epidemiol Infect* 2003; 103:301-312.
- Thomson DU, Loneragan GH, Thornton AB, Lechtenberg KF, Emery DA, Burkhardt DT, Nagaraja TG. Use of siderophore receptors and porin proteins-based vaccine to control the burden of *Escherichia coli* O157:H7 in feedlot cattle. *Foodborne Path Dis* 2009; 6:871-877.
- Tkalcic S, Brown CA, Harmon BG, Jain AV, Mueller EPO, Parks A, Jacobsen KL, Martin SA, Zhao T, Doyle MP. Effects of diet on rumen proliferation and fecal shedding of *Escherichia coli* O157:H7 in calves. *Appl Environ Microbiol* 2000; 70:5336-5342.
- Tournut J. Applications of probiotics to animal husbandry. *Rev Sci Tech Off Int Epiz* 1989; 8:551-566.
- Tutenel AV, Pierard D, Van Hoof J, De Zutter L. Molecular characterization of *Escherichia coli* O157 contamination routes in a cattle slaughterhouse. *J Food Prot* 2003; 66:1564-1569.
- USDA-FSIS. Recall Case Archive 2011. Accessed 3 Feb 2012.
http://www.fsis.usda.gov/fsis_recalls/Recall_Case_Archive_2011/index.asp
http://www.fsis.usda.gov/Fsis_Recalls/Open_Federal_Cases/index.asp
- USDA-FSIS. FSIS to extend implementation date on routine sampling program for non-O157:H7. Accessed 8 Feb 2012.
http://www.fsis.usda.gov/News_&_Events/Const_Update_020812/index.asp

- Van Baale MJ, Sargeant JM, Gnad DP, DeBey BM, Lechtenberg KF, Nagaraja TG. Effect of forage or grain diets with or without Monensin on ruminal persistence and fecal *Escherichia coli* O157:H7 in cattle. *Appl Environ Microbiol* 2004; 70:5336-5342.
- van der Waaij D, Berghuis de Vries JM, Lekkerkerk JEC. Colonization resistance of the digestive tract in conventional and antibiotic-treated mice. *J Hyg* 1971; 69:405-511.
- Viuda-Martos M, Ruiz-Navajas Y, FernAjndez-Lapez J, Perez-Alvarez J. Antibacterial activity of lemon (*Citrus lemon L.*), mandarin (*Citrus reticulate L.*), grapefruit (*Citrus paradise L.*) and orange (*Citrus sinensis L.*) essential oils. *J Food Safety* 2008; 28:567-576.
- Vogt RL, Dippold L. *Escherichia coli* O157:H7 outbreak associated with consumption of ground beef, June-July 2002. *Public Health Report* 2005; 120:174-178.
- Volanis M, Zoiopoulos P, Panagou E, Tzerkis C. Utilization of an ensiled citrus pulp mixture in the feeding of lactating dairy ewes. *Small Rum Res* 2006; 64:190-195.
- Wadolowski EA, Sung LM, Burris JA, Samuel JE, O'Brien AD. Acute renal tubular necrosis and death of mice orally infected with *Escherichia coli* strains that produce shiga-like toxin type II. *Infect Immun* 1991; 58:3959-3966.
- Wang G, Zhao T, Doyle MP. Fate of enterohemorrhagic *Escherichia coli* O157:H7 in bovine feces. *Appl Environ Microbiol* 1996; 62:2567-2570.
- Wetzel AN, LeJeune JT. Clonal dissemination of *Escherichia coli* O157:H7 subtypes among dairy farms in Northeast Ohio. *Appl Environ Microbiol* 2006; 72:2621-2626.
- Willard ST, Amstutz MD, Abraham E, Castaño JP, Leaumont DC, Faught WJ, Frawley LS. Simultaneous indirect activity measurements of GH and PRL genes in the same, living cell: identification of mammosomatotropes. *Am J Physiol* 1999; 277:E1150-E1153.
- Willard S, Ryan P, Bailey R, Lawrence M, Estill C, Gandy S, Lay JR,. Development of a novel paradigm for real-time monitoring of bacterial pathogenicity in swine. *J Anim Sci* 2002; 85(suppl. 1):31. (Abstr).
- Wolin MJ. Volatile fatty acids and the inhibition of *Escherichia coli* growth by rumen fluid. *Appl Microbiol* 1969; 17:83-87.

- Yang HE, Yang WZ, McKinnon JJ, Alexander TW, Li YL, McAllister TA. Survival of *Escherichia coli* O157:H7 in ruminal or fecal contents incubated with corn or wheat dried distillers' grain with solubles. *Can J Microbiol* 2010; 56:890-895.
- Zhao T, Tkalcic S, Doyle MO, Harmon BG, Brown CA, Zhao P. Pathogenicity of enterohemorrhagic *Escherichia coli* in neonatal calves and evaluation of fecal shedding by treatment with probiotic *Escherichia coli*. *J Food Prot* 2003; 66:924-930.
- Zhao H, Doyle TC, Wong RJ, Cao Y, Stevenson DK, Piwnica-Worms D, Contag CH. Characterization of coelenterazine analogs for measurements of *Renilla* luciferase activity in live cells and living animals. *Mol Imaging* 2004; 3:43-54.
- Zopf D, Roth S. Oligosaccharide anti-infective agents. *Lancet* 1996; 347:1017-1021.

CHAPTER III
SURVIVAL OF *ESCHERICHIA COLI* O157:H7 (ATCC 43888) TRANSFORMED
WITH EITHER THE PAK1-*LUX* OR PXEN-13 PLASMIDS IN BOVINE
MIXED RUMINAL AND FECAL MICROORGANISM FLUID

Abstract

The use of luminescent plasmids in bacteria may serve as a viable model for the real-time validation of various pre-harvest interventions on the colonization or shedding patterns of *Escherichia coli* O157:H7 within cattle. The objective of this study was to determine if the growth characteristics of *E. coli* O157:H7 (ATCC 43888) in mixed ruminal and fecal microorganism fluid cultures was altered when transformed with one of the two luminescent plasmids: pAK1-*lux* or pXEN-13. Transformants harboring the luminescent plasmids were compared to the non-transformed parental strain (WT) in mixed ruminal and fecal microorganisms fluid media for 6 h in triplicate (n = 3). The transformants and WT grew similarly in the presence of ampicillin (50 µg/mL). Within mixed ruminal microorganism fluid fermentations all transformants grew similarly through the 6 h study. The three transformants also grew similarly within mixed fecal microorganism fluid. The RLU (reflective light units; photon/pixel second) photonic emissions of each plasmid within ruminal fluid differed at 0 h ($P < 0.002$) and 2 h ($P < 0.02$) and within fecal fluid 0 h ($P < 0.009$) and 2 h ($P < 0.04$). The RLUs remained the same within rumen fluid at 4 h ($P < 0.22$) and 6 h ($P < 0.80$) and within fecal fluid at 4 h ($P < 0.06$) and 6 h ($P < 0.29$). Growth of *E. coli* O157:H7 transformed with the

bioluminescent plasmids was not altered in comparison to the WT, suggesting that both plasmids may serve as useful models for *in vivo* studies.

Introduction

One of the largest food safety threats facing the cattle industry today is the foodborne pathogenic bacterium *Escherichia coli* O157:H7 and other shiga toxin-producing *E. coli* (STEC; Gyles, 2007; Karmali et al., 2010). This bacterium is a natural member of the gastrointestinal microbial ecosystem of cattle; it does not cause illness in cattle, but can cause severe illness and even death in human consumers (Gyles, 2007; Karmali et al., 2010). Many intervention strategies aimed at reducing this critical pathogen have been implemented at the abattoir at a cost of more than \$2 billion (Kay, 2003; Koohmaraie et al., 2005; Arthur et al., 2009). *Escherichia coli* O157:H7 can also be transmitted directly by contact to farm workers and visitors, and it can be carried by water run-off to nearby water supplies or irrigated crops (Chapman et al., 2000). In 2010, STEC illnesses were estimated to cost \$1 billion dollars due to medical costs associated with foodborne illnesses and deaths in the United States (Scharff, 2010). Because of the link between this pathogen in live cattle and human illness, there has been an increased focus on the development of pre-harvest intervention strategies that reduce pathogenic bacteria in live food animals (Sargeant et al., 2007).

A limiting factor to pre-harvest intervention development has been the inability to physically track STEC or *E. coli* O157:H7 in a real-time fashion. Since contamination on meat carcasses could potentially originate from fecal material, punctured gastrointestinal contents and from feces on hides, a means of tracking of this pathogen through the use of biophotonics would provide a valuable model (Lohr, 1996). Detection of bioluminescent

cells is extremely sensitive when using an intensified charged-coupled device camera, with as few as 50 colony forming units being distinguishable (Siragusa et al., 1999). Bacteria transformed with luminescent plasmids that constitutively express the *luxCDABE* operon have been reported to have no alterations in growth kinetics compared to parental strains (Beyer and Bohm, 1996), and no significant alterations biochemically, serologically, or structurally (Chen and Griffiths, 1996).

The ability to track bioluminescent pathogens *in vivo* through the gastrointestinal tract has previously been successful in pigs and mice (Willard et al., 2002; Burns-Guydish et al., 2005; Moulton et al., 2009a). Employing bioluminescence in live animal models will facilitate tracking of pathogens (Contag et al., 1997) and can indicate the physical location of these pathogens (Siragusa et al., 1999) which will increase researchers' understanding of pathogenic processes (Contag et al., 1995). However, limited information is available for the capability of this technology for monitoring *E. coli* colonization and shedding *in vivo*. In particular, it is not known whether bioluminescent plasmids can be retained within the stressful, competitive environment of the gastrointestinal tract. Therefore, the objectives of this study were to analyze the parental strain *E. coli* O157:H7 ATCC 43888 wild type (WT) transformed separately with two different bioluminescent plasmids, pAK1-*lux* and pXEN-13, in order to determine whether the growth of *E. coli* O157:H7 was altered in bovine mixed ruminal microorganism fluid or fecal microorganism fluid media in relation to the non-transformed parental strain. The present study is a preliminary step in the validation of the use of bioluminescent *E. coli* O157:H7 as a novel real-time model for foodborne pathogen contamination.

Materials and Methods

Bacterial Cultivation Conditions

Escherichia coli ATCC 43888 is an O157:H7 serotype that does not possess the shiga toxin I or shiga toxin II genes. This bacterium was routinely cultured on Tryptic Soy Agar (TSA; Difco Co., Corpus Christi, TX) at 37°C. The transformation of *E. coli* O157:H7 with the plasmid pXEN-13 (Caliper Life Sciences, Hopkinton, MA), cells were washed three times with 10% glycerol and electroporated with 50 µg of the pXEN-13 plasmid (Sambrook and Russell, 2001). The *E. coli* O157:H7 ATCC 43888 was transformed with pAK1-*lux* as previously described by Moulton et al., 2008. Both transformed *E. coli* O157:H7 were selected for growth on 50 µg/mL of ampicillin (AMP) at 37°C; bioluminescence was confirmed by analyzing transformed strains with a Berthold/NightOwl camera equipped with the WinLight 32 software, version 2.51.11901 (Berthold Technologies; Oak Ridge, TN). The 43888 strain was passed through a series of increasing concentrations of novobiocin until the cells were made resistant to 10 µg/ml novobiocin.

Pure Culture Growth

Overnight cultures of the parental strain 43888, 43888 containing pAK1-*lux*, and 43888 containing pXEN-13 were cultured in triplicate at 37°C with constant agitation (140 rpm) in TSB supplemented with 50 µg/mL AMP or 10 µg/mL novobiocin. Overnight cultures (2 mL) were pelleted for 2 min at 10,000 x *g*, washed once with 2 mL of 1X phosphate buffer solution (PBS), and resuspended in 2 mL of TSB in order to remove the antibiotics. Cultures were then diluted 1:100 in TSB supplemented with 50 µg/mL AMP or 10 µg/mL novobiocin, and 200 µL aliquots were transferred, in duplicate,

to a 96-well microtiter plate. The OD₆₀₀ for each culture grown in both conditions at 37°C was measured hourly for 24 h to verify that the plasmids did not slow growth rate, using a BioTek (Winooski, VT, USA) Synergy HT multi-mode microplate reader and analyzed with Gen5 data analysis software. Maximum specific growth rate was measured during exponential phase, while OD₆₀₀ was less than 0.6 OD₆₀₀ units and calculated according to Callaway and Russell (1999).

Mixed Ruminant and Fecal Microorganism Contents Collection

Rumen contents (1000 mL) were collected from the rumen ventral sac of a 362 kg cannulated steer at the Henry Leveck Animal Research Center at Mississippi State University. Fecal contents (6000 mL) were collected rectally from six Holstein cows at the Joe Bearden Dairy Research Center at Mississippi State University. Particles were separated from the ruminal and fecal fluid by passing contents through nylon paint strainers as previously described by Leyendecker et al., 2004. After separation, rumen and fecal fluids were incubated for 30 min at 37°C, which resulted in the formation of three visible layers. The middle layer of both ruminal and fecal fluid were removed and utilized for the foundation of media. The base medium utilized for both rumen and fecal fluid contained (per liter): 6.0 g KH₂PO₄, 6.0 g KH₂PO₄, 12.0 g (NH₄)₂SO₄, 12.0 g NaCl, 2.5 g MgSO₄•7 H₂O, 1.6 g CaCl₂•2H₂O, 0.04 g cysteine HCl; base medium was sterilized by autoclaving. Then 33.0 mL of an 8% solution of Na₂CO₃ and 333 mL of ruminal fluid or fecal fluid were added and homogenized by mixing. The pH was adjusted to 6.5 with 1 M of NaOH solution and bubbled with CO₂. Both fully prepared media (Cotta and Russell, 1982) were incubated in an orbital shaker at 140 rpm at 37°C for 12 h.

Mixed Ruminant and Fecal Microorganism Contents Fermentations

Fresh overnight cultures, in triplicate prepared as described above, of the parental strain 43888, 43888 containing pAK1-*lux*, and 43888 containing pXEN-13 grown in TSB supplemented with the appropriate antibiotic were then diluted 1:100 in 2 mL of TSB and allowed to incubate with agitation (140 rpm) at 37°C until mid-log phase ($OD_{600} \sim 0.50$; Broadway, 2011) was achieved. Cultures were then split and aliquotted; one tube for ruminal fluid and one for fecal fluid, pelleted by centrifugation at 10,000 x *g* for 2 min, and resuspended in 5 mL of either media containing ruminal or fecal fluid medium. After inoculation, cultures were incubated with constant agitation (140 rpm) at 37°C for 6 h. At each 120 min interval, four 100 mL aliquots of each culture were transferred to a 96-well plate (Costar® Black Sterile Polystyrene; Corning Incorporated, Corning, NY) and subsequently imaged. Photonic emissions were recorded following a 2 min acquisition phase and were measured based on reflective light units [RLU; photons/pixel per second (ph/pix s)]. Additionally, aliquots of each culture were acquired at 0 h and continuously at 120 min intervals for 6 h, diluted in 1X PBS, and plated onto the appropriate selective media: parental strain 43888 on TSA+ 10 mg/mL novobiocin, 43888 containing pAK1-*lux* or pXEN-13 on TSA+50 mg/mL AMP. Plates were then incubated at 37°C for 12 h prior to analysis, viable plate counts were obtained. Plates were also imaged using a Berthold/NightOwl camera to validate luminescence.

Statistical Analysis

Data were analyzed as a completely randomized design with repeated measures using PROC MIXED in SAS (SAS Inst. Inc., Version 9.2; Cary, NC). Experimental unit was defined as tube, and significance was declared at $P < 0.05$. Pair-wise differences

among least squares means at various sample times were evaluated with the PDIFF statement.

Results

Growth of *E. coli* 43888 containing pAK1-*lux* or pXEN-13

To determine whether pAK1-*lux* and pXEN-13 plasmids exhibited similar growth rates, growth was monitored over a 24 h period using a BioTek Synergy HT plate reader (Winooski, VT, USA) [Figure 3.1]. The pAK1-*lux* (1.05 h^{-1}) and the pXEN-13 (1.02 h^{-1}) containing-transformants both exhibited similar maximum specific growth rates as the parental strain 43888 (0.99 h^{-1} ; Figure 3.1A). No differences were evident between the growth rates of pXEN-13 and the parental strain. Growth of both the pAK1-*lux* and pXEN-13 containing-transformants were also examined in the presence of the selective pressure of ampicillin (Figure 3.2B). The growth patterns of both transformants were similar, pAK1-*lux* (1.41 h^{-1}) and pXEN-13 (1.44 h^{-1}) and with no differences observed.

Growth in Ruminal Fluid Fermentations

To determine whether differences were evident in the ability of the two plasmids to be retained within the rumen; both strains containing either pAK1-*lux* or pXEN-13 were grown in the presence of freshly prepared mixed rumen microorganism fluid medium. The parental strain was also grown under these conditions to determine whether the presence of the plasmids would alter the ability of *E. coli* O157:H7 to survive. The *E. coli* O157:H7 and the two transformants grew similarly within mixed ruminal microorganism fluid (Figure 3.2). Growth was monitored by viable plate counts and populations were recorded in \log_{10} (CFU/mL) populations. There was no difference ($P > 0.18$) observed within populations between the WT and the pXEN-13 plasmid throughout

the study. The only difference observed within the study was at 4 h, when pAK1-*lux* had greater populations than pXEN-13 ($P < 0.004$), but was not different from the WT ($P > 0.10$).

Growth in Fecal Fluid Fermentations

To determine whether the presence of either plasmid would affect the ability of *E. coli* O157:H7 to survive within mixed fecal microorganism fluid, strains containing either pAK1-*lux* or pXEN-13 were grown in the presence of freshly prepared fecal fluid medium and compared to growth of the non-transformed parental strain. The WT and the two transformants grew similarly within mixed fecal microorganism fluid fermentations (Figure 3.3). There was no difference ($P > 0.14$) observed within populations between the WT and pXEN-13 for the duration of the study. The only difference observed was at 6 h when pXEN-13 suggested greater populations than pAK1-*lux* ($P < 0.01$) but was not different than the WT ($P < 0.18$).

Comparison of Photonic Emissions by pAK1-*lux* and pXEN-13

To determine whether photonic light emission would be sustained by both pAK1-*lux* and pXEN-13, RLU photonic emissions were monitored over a 6 h period in both mixed rumen and fecal microorganism fluid media. Photonic emissions gradually decreased from 0 h for both plasmids within ruminal fluid, indicating that there was a time effect ($P < 0.001$). The RLU values are measured in photons/pixel second (ph/pix s). The RLU values for pAK1-*lux* ranged from 16225.97 ph/pix s to 880.21 ph/pix s (data not reported in tabular form). The pXEN-13 plasmid had the greatest RLU values at 0 h at 292.28 ph/pix s and the lowest at 6 h 86.25 ph/pix s. The pAK1-*lux* plasmid had greater RLU values and was statistically different at 0 h ($P < 0.002$) and 2 h ($P < 0.02$)

and as photonic emissions decreased over time both plasmids emitted the similar values at 4 h ($P < 0.22$) and 6 h ($P < 0.80$), suggesting that there was only a tendency ($P < 0.07$) for a transformant effect. There was a transformant by time effect ($P < 0.001$) because both time and transformant affect photonic emissions.

Within mixed fecal microorganisms fluid media both plasmid RLU values decreased from 0 h, indicating that time impacted photonic emissions ($P < 0.001$). The pAK1-*lux* plasmid ranged from 5513.65 ph/pix s to 1404.42 ph/pix s (data not reported in tabular form). The pXEN-13 plasmid had decreased RLU values with a range from 112.92 ph/pix s to 34.61 ph/pix s. The two plasmids were different from 0 h ($P < 0.009$) and 2 h ($P < 0.04$), with a tendency to be different at 4 h ($P < 0.06$). At 6 h the plasmids were not different ($P < 0.29$), further suggesting that a tendency for transformant ($P < 0.06$) to impact photonic emissions, while indicating a transformant by time interaction ($P < 0.001$).

Discussion

To examine the application of bioluminescent technology in a large animal study, we analyzed the growth and luminescence of *E. coli* 43888 containing either the pAK1-*lux* or the pXEN-13 plasmid. Bioluminescence studies have suggested correlations with *in vitro* enzyme activity; the amount of luciferase protein and *lux* mRNA generated directly correlates with cell viability (Close et al., 2010). The two plasmids utilized in this study were selected on the basis of harboring the *luxCDABE* operon. The pAK1-*lux* plasmid was developed as a wide range host plasmid for gram-negative bacteria (Karsi et al., 2006). Research with this plasmid has suggested stability and luminescence, without antibiotic pressure for at least 8 d; while decreasing the percentage of photonic emissions

from 0 d (Moulton et al., 2008). The pXEN-13 plasmid carries the original *Photorhabdus luminescens* operon for production of luminescence in gram-negative bacteria (Harms et al., 2009). The pXEN-13 plasmid has been introduced into *E. coli* strains previously and emitted photonic emissions within the mouse model (Harms et al., 2009). The WT containing either pAK1-*lux* or pXEN-13 exhibited similar growth rates under antibiotics pressure.

Both *luxCDABE* containing plasmids grew similarly and no differences were noted in the growth of the plasmid-equipped bacteria in the presence of AMP, indicating that both plasmids grow similarly to the WT parental strain in the absence of AMP, and similarly under AMP pressure. This suggests that even within a live animal model without antibiotic pressure, both transformants will function similarly to the WT.

Many studies have demonstrated that *E. coli* O157:H7 can be cultured from the rumen, but does not grow rapidly or extensively in the rumen (Grauke et al., 2002). To further investigate the growth and stability of WT and the two bioluminescently transformed *E. coli* O157:H7, pAK1-*lux* and pXEN-13 were inoculated and grown in mixed microbial ruminal fluid microorganism fermentations. There was no difference observed in the two transformants compared with the controls when grown in mixed microbial ruminal fluid fermentations, thus suggesting that both transformants can survive the harsh rumen conditions and grow in a fashion similar to the WT.

Escherichia coli O157:H7 can contaminate beef carcasses via fecal spread from hide or spilled digesta and it has been suggested that 30% of cattle harbor *E. coli* O157:H7 or other EHEC in feces, depending on season (Elder et al., 2000; Callaway et al., 2003). Transformants and WT grew similarly in the presence of a mixed fecal microorganism fluid environment throughout the duration of the study. This suggests

that transformants functioned similarly within the fecal fluid environment and are capable of surviving the competitive fecal environment.

Transforming bacteria with the bioluminescent plasmids has been suggested to be a successful means of tracking bacteria in real-time (Moulton et al., 2008; Moulton et al., 2009b; Curbelo et al., 2010). The present data suggest that the pAK1-*lux* and pXEN-13 transformants are stable and can persist in rumen and fecal-like environments.

Furthermore, while grown in these harsh, competitive environments, these transformants still display luminescence and could be a useful method for pathogen tracking in real-world environments (Moulton et al., 2008). The use of luciferase reporter genes in real-time imaging can aid in pre-harvest pathogen intervention *in vivo* by reducing the number of animals needed for use in studies, as well as by providing the physical location of pathogen contamination routes (Ryan et al., 2005).

Conclusions

Transformation of *E. coli* O157:H7 with the pAK1-*lux* gene cassette or the pXEN-13 gene cassette did not alter the growth of the transformants compared to the WT (ATCC 43888) within mixed ruminal or mixed fecal microorganism incubations. Our results suggest that both plasmids could be implemented for a model for pre-harvest pathogen intervention in ruminant animals.




Figure 3.1 The OD₆₀₀ of cultures of the wild type (WT) *Escherichia coli* O157:H7 ATCC 43888 (◆) and the two bioluminescently transformed variants, 43888 containing the plasmid pXEN-13 (▲) or pAK1-*lux* (■). Cultures were grown in the absence of (A) and presence (B) of ampicillin (50 mg/mL) to provide selective pressure




Figure 3.2 The least square means of *Escherichia coli* O157:H7 (ATCC 43888) populations (\log_{10} CFU/mL) in bovine mixed ruminal microorganism incubations. Growth was monitored by viable plate counts with the parental strain (◆; WT) or 43888 containing the plasmid pAK1-*lux* (▲; PAK) or pXEN-13 (■; XEN) in bovine mixed ruminal fluid for 6 h. Values represent the \log_{10} CFU/mL values from three independent replicates

^{x,y} Lsmeans lacking a common subscript within column differ ($P < 0.05$)




Figure 3.3 The least square means of *Escherichia coli* O157:H7 (ATCC 43888) populations (\log_{10} CFU/mL) in bovine mixed fecal microorganism incubations. Growth was monitored by viable plate counts with the parental strain (◆; WT) or 43888 containing the plasmid pAK1-*lux* (▲; PAK) or pXEN-13 (■; XEN) in bovine mixed fecal fluid for 6 h. Values represent the \log_{10} CFU/mL values from three independent replicates

^{x,y} Lsmeans lacking a common subscript within column differ ($P < 0.05$)

Literature cited

- Arthur TM, Keen JE, Bosilevac JM, Brichta-Harhay DM, Kalchayanand N, Shackelford SD, Wheeler TL, Nou X, Koohmaraie M. Longitudinal study of *Escherichia coli* O157:H7 in a beef cattle feedlot and role of high-level shedders in hide contamination. *Appl Environ Microbiol* 2009; 75:6515-6523.
- Beyer W, Böhm R. Labeling *Salmonella* live vaccine strains with the *lux* operon from *Vibrio fischeri* improves their detection and decimation from wild type. *Microbiol Res* 1996; 151:407-419.
- Broadway PR. Stability of biophotonic *Escherichia coli* O157:H7 and its effectiveness as a validation tool in beef purge sampling methodologies. Masters Theses; 2011 Mississippi State Univ., Starkville.
- Burns-Guydish SM, Olomu IN, Zhao H, Wong RJ, Stevenson DK, Contag CH. Monitoring age-related susceptibility of young mice to oral *Salmonella enterica* serovar *Typhimurium* infection using *in vivo* murine model. *Pediatr Res* 2005; 58:153-158.
- Callaway TR, Russell JB. Selection of a highly monensin-resistant *Prevotella bryantii* subpopulation with altered outer membrane characteristics. *Appl Environ Microbiol* 1999; 65:4753-4759.
- Callaway TR, Edrington TS, Anderson RC, Genovese KJ, Poole TL, Elder RO, Byrd JA, Bischoff KM, Nisbet DJ. *Escherichia coli* O157:H7 populations in sheep can be reduced by chlorate supplementation. *J Food Prot* 2003; 66:194-199.
- Chapman PA, Cornell J, Green C. Infection with verocytotoxin-producing *Escherichia coli* O157 during a visit to an inner city open farm. *Epidemiol Infect* 2000; 125:531-536.
- Chen J, Griffiths MW. *Salmonella* detection in eggs using *lux+* bacteriophages. *J Food Prot* 1996; 59:908-914.
- Close DM, Patterson SS, Ripp S, Baek SJ, Sanseverino J, Saylor GS. Autonomous bioluminescent expression of the bacterial luciferase gene cassette (*lux*) in a mammalian cell line. *PLoS ONE* 2010; 5:e12441.
- Contag CH, Contag PR, Mullins JI, Spilman SD, Stevenson DK, Benaron DA. Photonic detection of bacterial pathogens in living hosts. *Mol Microbiol* 1995; 18:593-603.

- Contag CH, Spillman SD, Contag PR, Oshiro M, Eames B, Dennery P, Stevenson DK, Benaron DA. Visualizing gene expression in living mammals using a bioluminescent reporter. *Photchem Pathobiol* 1997; 66:523-531.
- Cotta MA, Russell JB. Effect of peptides and amino acids on efficiency of rumen bacterial protein synthesis in continuous culture. *J Dairy Sci* 1982; 65:226-234.
- Curbelo J, Moulton K, Willard S. Photonic characteristics and *ex vivo* imaging of *Escherichia coli* Xen-14 within the bovine reproductive tract. *Therio* 2010; 73:48-55.
- Elder RO, Keen JE, Siragusa GR, Barkocy-Gallagher GA, Koohmaraie M, Laegreid WW. Correlation of enterohaemorrhagic *Escherichia coli* O157 prevalence in feces, hides and carcasses of beef cattle during processing. *Proc Natl Acad Sci USA* 2000; 97:2999-3003.
- Grauke, LJ, Kudva IT, Yoon JW, Hunt CW, Williams CJ, Hovde CJ. Gastrointestinal tract location of *Escherichia coli* O157:H7 in ruminants. *Appl Environ Microbiol* 2002; 68:2269-2277.
- Gyles CL. Shiga toxin-producing *Escherichia coli*: An overview. *J Anim Sci* 2007; 85:E45-E62.
- Harms JS, Durward MA, Magnani DM, Splitter GA. Evaluation of recombinant invasive, non-pathogenic *Escherichia coli* as a vaccine vector against the intracellular pathogen, *Brucella*. *J Immun Ther Vaccines* 2009; 7:1.
- Karmali MA, Gannon V, Sargeant JM. Verocytotoxin-producing *Escherichia coli* (VTEC). *Vet Microbiol* 2010; 140:360-370.
- Karsi A, Menanteau-Ledouble S, Lawrence ML. Development of bioluminescent *Edwardsiella ictaluri* for noninvasive disease monitoring. *FEMS Microbiol Lett* 2006; 260:216-223.
- Kay S. *E. coli* O157:H7: The cost during the last 10 years. *Meat and Poultry News* 2003; 3:26-34.
- Koohmaraie M, Arthur TM, Bosilevac JM, Guerini M, Shackelford SD, Wheeler TL. Post-harvest interventions to reduce/eliminate pathogens in beef. *Meat Sci* 2005; 71:79-91.
- Leyendecker SA, Callaway TR, Anderson RC, Nisbet DJ. Technical note on a much simplified method for collecting ruminal fluid using a nylon paint strainer. *J Sci Food Agric* 2004; 84:387-389.

- Lohr JA. Beef carcass microbial contamination-post slaughter numbers of bacteria, sources of contamination and variability. Proceedings of 49th Annual Reciprocal Meats Conference, Provo, Utah 1996; 132-137.
- Moulton K, Ryan P, Christiansen D, Hopper R, Klausner C, Bennett W, Rodta-Palenik, Willard ST. *Ex vivo* bioluminescence imaging of late gestation ewes following intrauterine inoculation with *lux*-modified *Escherichia coli*. *Comp Immunol Microbiol Infect Dis* 2008; 32:429-438.
- Moulton K, Ryan P, Lay D, Willard S. Postmortem photonic imaging of *lux*-modified *Salmonella typhimurium* within the gastrointestinal tract of swine after oral inoculation *in vivo*. *J Anim Sci* 2009a; 87:2239-2244.
- Moulton K, Ryan P, Lay D, Willard S. Photonic plasmid stability of transformed *Salmonella Typhimurium*: A comparison of three unique plasmids. *BMC Microbiology* 2009b; 9:152.
- Ryan P, Youngblood RC, Harvill J, Willard ST. Photonic monitoring in real time of vascular endothelial growth factor receptor 2 gene expression under relaxin-induced conditions in a novel murine wound model. *Ann NY Acad Sci* 2005; 1041:398-414.
- Sambrook J, Russell DW. *Molecular cloning: a laboratory manual*, 3rd edition. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, 2001.
- Sargeant JM, Amezcua MR, Rajic A, Waddell L. Pre-harvest interventions to reduce the shedding of *E. coli* O157 in the faeces of weaned domestic ruminants: a systematic review. *Zoonos Pub Health* 2007; 54:260-277.
- Scharff, RL 2010. Health-related costs from foodborne illness in the United States. 3 May, Accessed 2010.
<http://www.producesafetyproject.org/admin/assets/files/Health-Related-Foodborne-Illness-Costs-Report.pdf-1.pdf>.
- Siragusa GR, Nawotka K, Spilman SD, Contag PR, Contag CH. Real-time monitoring of *Escherichia coli* O157:H7 adherence to beef carcass surface tissues with a bioluminescent reporter. *Appl Environ Microbiol* 1999; 65:1738-1745.
- Willard S, Ryan P, Bailey R, Lawrence M, Estill C, Gandy S, Lay JR. Development of a novel paradigm for real-time monitoring of bacterial pathogenicity in swine. *J Anim Sci* 2002; 85(suppl. 1):31. (Abstr).

CHAPTER IV

EFFECTS OF CITRUS BY-PRODUCTS ON GROWTH OF O157:H7 AND NON-O157 *ESCHERICHIA COLI* SEROTYPES WITHIN BOVINE MIXED RUMINAL MICROORGANISM FLUID

Abstract

Citrus by-products (CBP) are utilized in citrus regions of the United States as a low cost, highly palatability and nutritional component of cattle diets. Researchers have suggested that inclusion of CBP both *in vivo* and *in vitro* can inhibit the growth of both *Escherichia coli* O157:H7 and *Salmonella* (Nannapaneni et al., 2008; Collier et al., 2010). The objective of this study was to examine the effects *in vitro* that varying concentrations of CBP in the powdered or pelleted variety impose on enterohemorrhagic *Escherichia coli* (EHEC) serotypes; O26:H11, O103:H8, O111:H8, O145:H28, O157:H7 and O157:H7 $\Delta stx_1 stx_2$ in mixed ruminal microorganism media. Mixed ruminal microorganism fluid media were supplemented with either powdered or pelleted CBP at concentrations of 0%, 5%, 10%, 20% and EHEC growth (\log_{10} CFU/mL) were evaluated at 0, 2, 4, 6 h. The O26:H11, O111:H8, O145:H28 and O157:H7 serotypes did not exhibit a change in populations throughout both CBP varieties. The O103:H8 serotype displayed an approximate 1 \log_{10} reduction in O103:H8 populations at 5% powdered CBP over 6 h. There was an approximate 1.5 \log_{10} reduction in O157:H7 $\Delta stx_1 stx_2$ populations at 10% CBP and an approximate 5 \log_{10} reduction observed in O157:H7 $\Delta stx_1 stx_2$ populations at 20% powdered CBP over the 6 h study. There was an

approximate 1 log₁₀ reduction in O103:H8 populations when supplemented with 20% pelleted CBP. These results suggest that the serotypes of EHEC grow differently within mixed ruminal microorganism fluid media when supplemented with CBP in the powdered or pelleted variety. Further research is needed to indicate the affect that CBP have on non-O157 EHEC serotypes.

Introduction

Enterohemorrhagic *Escherichia coli* (EHEC) are capable of naturally colonizing within the gastrointestinal tract of cattle, without causing illness (Wells et al., 1991). Human consumption of products contaminated with EHEC can cause illness and more severely hemorrhagic colitis and hemorrhagic uremic syndrome (Griffin and Tauxe, 1991; Stanford et al., 2005). The most notorious EHEC within the meat industry has been *E.coli* O157:H7. Due to increased surveillance and pre- and post- harvest intervention, the occurrence of O157:H7 infections have been reduced to ≤ 1 case per 100,000 people (CDC, 2011a). However, there now appears to be an increase in the occurrence of foodborne outbreaks due to non-O157 EHEC (CDC, 2011a). According to the Center for Disease Control (CDC) an estimated 265,000 cases of EHEC were reported a year; of these, approximately 67% are attributed to non-O157 EHEC (CDC, 2011b). With increased concerns related to the prevalence of non-O157 outbreaks the USDA-FSIS will be labeling non-O157 EHEC serotypes O26, O45, O103, O111, O121 and O145 as adulterants in fresh non-intact beef products (USDA-FSIS, 2012).

The production of citrus for various food and non-food products generates by-products such as the pulp and peel from citrus fruit. These citrus by-products (CBP) have been utilized by dairy and beef cattle producers in regions of the United States as an

inexpensive nutritionally dense feed source (Callaway et al., 2008). The incorporation of CBP into diets for cattle also may aid in the reduction of foodborne pathogens due to antimicrobial aspects of the by-products. Citrus products and by-products contain essential oils that possess antimicrobial activities that are able to damage the cell wall due to lipophilic properties. The lipophilic properties of essential oils are able to permeabilize the cell wall within gram-negative bacteria (Bakkali et al., 2006). The change in the fluidity of the membranes due to the permeabilization, allows essential oils to coagulate in the cytoplasm (Gustafson et al., 1998), depleting ATP (Di Pasqua et al., 2006) and resulting in lysis of the cell (Kim et al., 2005; Fisher and Phillips, 2006; Gill and Holley, 2006; Oussalah et al., 2006; Di Pasqua et al., 2007). Research pertaining to essential oils within CBP has indicated severe damage to *E. coli* serotypes (Kim et al., 1995; Dusan et al., 2006). The rumen and intestinal gram-negative microbial populations of cattle can be altered due to this antimicrobial activity within cattle (Nam et al., 2006). Since CBP contain antimicrobial properties and are readily available at low costs within citrus-producing areas and has nutritional value, it is being investigated as a potential pre-harvest pathogen intervention strategy to reduce EHEC concentrations within the gastrointestinal tract of cattle. Therefore, the objective of this study was to examine the effects of powdered and pelleted citrus by-products have on growth of the EHEC serotypes: O26:H11, O103:H8, O111:H8, O145:H28, O157:H7 and O157:H7 $\Delta stx_1 stx_2$ in mixed ruminal microorganism media.

Materials and Methods

Ruminal Fluid Collection and Medium Preparation

Ruminal contents (1000 mL) were collected from the rumen ventral sac of a 544 kg cannulated steer at the Henry Leveck Animal Research Center at Mississippi State University. Rumen particles were separated from the ruminal fluid by passing contents through nylon paint strainers as previously described by Leyendecker et al., 2004. After separation, rumen fluid was incubated for 30 min at 37°C, to allow the fluid to separate into three distinct layers. The middle layer of the rumen fluid was extracted and utilized for the mixed ruminal microorganism fluid. The medium utilized for the mixed ruminal microorganism fluid contained (per liter): 6.0 g KH_2PO_4 , 6.0 g KH_2PO_4 , 12.0 g $(\text{NH}_4)_2\text{SO}_4$, 12.0 g NaCl, 2.5 g $\text{MgSO}_4 \cdot 7 \text{H}_2\text{O}$, 1.6 g $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 0.04 g cysteine HCl; base medium was sterilized by autoclaving (Cotta and Russell, 1982). To the base medium, 33.0 mL of an 8% solution of Na_2CO_3 and 333 mL of ruminal fluid were added and homogenized by mixing. The pH was adjusted to 6.5 with 1 M of NaOH solution, and bubbled with CO_2 . The fully prepared media was incubated in an orbital shaker at 140 rpm at 37°C for 12 h.

Bacterial Serotypes and Cultivation Conditions

Six serotypes of *E. coli* included in this study were purchased from the American Type Culture Collection (ATCC): O157:H7 (ATCC 43895), O103:k:H8 (ATCC 23982), O145:H28 (BAA-2129), O26:H11 (BAA-1653), O111:H8 (BAA-179) and O157:H7 $\Delta\text{stx}_1\text{stx}_2$ (ATCC 43888). All *E. coli* serotypes were routinely cultured in Luria-Bertani medium (LB; Difco Co.; Corpus Christi, TX) at 37°C. All serotypes were transformed with the bioluminescent pXEN-13 plasmid (Caliper Life Sciences; Hopkinton, MA), as a

marker for means of antibiotic selection for identification following incubation. For transformation with the pXEN-13 plasmid all serotypes were made competent by washing mid-log cultures four times with ice cold 10% glycerol. Competent cells were then transformed with pXEN-13 by electroporation and cultured in LB supplemented with 100 mg/mL of ampicillin (AMP) using standard techniques (Sambrook and Russell, 2001).

Citrus By-Product Trial

Isolates from fresh streaks of each serotype transformed with pXEN-13 on LB agar supplemented with 100 µg/mL AMP were used to inoculate a 5 mL starter culture in LB broth+AMP at 16 h at 37°C. Cultures were then diluted 1:100 in LB broth+AMP and allowed to grow to mid-log phase, after which cultures were pelleted, residual medium was removed, and cells were resuspended in an equal volume of mixed ruminal microorganism fluid media supplemented with 0, 5, 10 or 20% CBP. Cultures were incubated at 37°C at 140 rpm. Aliquots were removed at various times, diluted in 1X phosphate buffered saline (PBS), plated onto LB agar supplemented with 100 µg/mL AMP, and incubated overnight at 37°C. The pH values were measured from each serotype at each time interval at the various CBP concentrations recorded.

Statistical Analysis

Data were analyzed as a completely randomized design with repeated measures using PROC MIXED in SAS (SAS Inst. Inc., Version 9.2; Cary, NC). Experimental unit was defined as tube, and significance was declared at $P < 0.05$. Pair-wise differences among least squares means at various sample times were evaluated with the PDIFF statement.

Results

Citrus By-Products pH Values

The pH values of each CBP variety at 0%, 5%, 10% and 20% at each time interval were recorded. The pH was adjusted to 6.5 at -2 h prior to the addition of CBP in either variety. From -2 h to 6 h, pH increased within all serotypes from 6.5 to 7.5 for 0% CBP. At 0 h the pH was 5.5 and decreased to 4.0, in the 5%, 10% and 20% CBP throughout the study for both varieties. The pH values of the powdered CBP decreased according to the increasing concentrations over the time intervals. The pelleted CBP variety indicated a similar trend, with pH values decreasing with the increase in pelleted CBP concentrations across the 6 h study.

Powdered Citrus By-Product

The growth of the various EHEC populations (\log_{10} CFU/mL) within mixed ruminal microorganism fluid media indicated differences within serotypes (Table 4.1). The O26:H11 and O145:H28 serotypes grew similar ($P < 0.11$) within the powdered CBP. The only difference observed in O26:H11 serotype's decreased ($P < 0.006$) populations at 4 h supplemented with 20% powdered CBP throughout the study. The O103:H8 serotype exhibited approximately a 1 \log_{10} reduction in populations over the 6 h study, when mixed ruminal microorganism media was supplemented with 5% powdered CBP. The O157:H7 $\Delta stx_1 stx_2$ and O157:H7 serotypes had decreased ($P < 0.04$ and $P < 0.05$, respectively) populations in comparison to the other serotypes at 0 h. Although both O157 serotypes tended to grow similarly ($P < 0.06$) throughout 4 h, there was a difference observed at 6 h when O157:H7 $\Delta stx_1 stx_2$ had greater decreases ($P < 0.03$) in populations in comparison to O157:H7. When supplemented with 10% powdered CBP

the O157:H7 $\Delta stx_1 stx_2$ serotype displayed approximately a 1.5 log₁₀ reduction in populations, and at 20% powdered CBP there was approximately a 5 log₁₀ reduction in populations over the 6 h course of the study.

Pelleted Citrus By-Product

The growth of the EHEC populations within mixed ruminal microorganism fluid media indicated differences between the various serotypes (Table 4.2). The pelleted CBP tended to have no change ($P < 0.07$) in populations of O145:H28 from 0 h to 6 h. While O157:H7 had decreased ($P < 0.02$) at 0 h, there were no differences in populations observed between the O103:H8 and O157:H7 ($P < 0.11$) the remainder of the study. Populations of O103:H8 were decreased ($P < 0.02$) at 0 h, while populations tended to be similar ($P < 0.06$) to O111:H8 throughout the study. When mixed ruminal microorganism media was supplemented with 20% pelleted CBP there was approximately a 1 log₁₀ reduction observed over the 6 h study. The O26:H11 serotype populations decreased ($P < 0.05$) throughout the study, exhibiting the least populations at 4 h. *Escherichia coli* O157:H7 $\Delta stx_1 stx_2$ exhibited the most decreased ($P < 0.05$) EHEC populations at 0 h and throughout the study.

Discussion

A study conducted by Free et al. (2012) utilized the same serotypes within this study O26:H11, O103:H8, O111:H8, O145:H28, O157:H7 and O157:H7 $\Delta stx_1 stx_2$. The results of this study suggested that all serotypes were capable of growing with mixed ruminal microorganism fluid media; with decreased concentrations of serotypes O103:H8 and O145:H28 after 24 h compared to O157:H7 (Free et al., 2012). These data suggests the possibility that not all non-O157 serotypes function similarly within cattle.

The O157:H7 $\Delta stx_1 stx_2$ data indicated decreased populations within the mixed rumen microorganism fluid media supplemented with powdered CBP, while O103:H8 indicated decreased populations within both varieties of CBP. These results are in accordance with previous studies that have suggested a decrease in O157:H7 populations using other varieties of CBP. Callaway et al. (2008) supplemented mixed ruminal microorganism fluid media with 0%, 0.5%, 1% and 2% dried orange pulp and *E. coli* O157:H7 populations decreased according to increasing concentrations. While another Callaway et al. (2011) study supplemented sheep rations with 0%, 5%, 10% orange peel and reduced *E. coli* O157:H7 populations *in vivo*.

The CBP was added to the mixed ruminal microorganism fluid media 2 h before the serotypes were added to the mixture (0 h). Although a decrease in O103:H8 and O157:H7 $\Delta stx_1 stx_2$ populations were observed, other EHEC populations were not affected. Given that this study was only conducted for 6 h, the effects of CBP within the mixed ruminal microorganism fluid media and EHEC serotypes may not have been fully observed within the short time frame. Other studies indicate that CBP decrease *E. coli* O157:H7 populations from 24 h to 72 h (Callaway et al., 2008; Callaway et al., 2011). This study was only conducted for 6 h; an increased duration of the study could have been more beneficial to observe the effects of CBP on the various serotypes.

The essential oils within the CBP can permeabilize the bacterial cell walls and cytoplasm, leading to bacterial lysis, thus shifting the rumen environment leading to an increase in short-chain fatty acids while decreasing the pH. The acidic environment creates less favorable conditions for microbial populations to survive and replicate within, thus decreasing the possibility of *E. coli* O157:H7 populations. Although our research has reported a decrease in pH values with increasing CBP concentrations and an

observed decrease in O103:H8 and O157:H7 $\Delta stx_1 stx_2$ populations, this same trend was not observed within other EHEC serotypes. Further research is needed to determine how the various EHEC serotypes affect *E. coli* populations within mixed ruminal microorganism fluid media when supplemented with CBP.

Conclusion

Enterohemorrhagic *Escherichia coli* are foodborne pathogens that can survive within the gastrointestinal tract of cattle without imposing illness. Essential oils are the main component of citrus by-products and have been suggested to contain antimicrobial properties. Although decreased populations were observed within the O103:H8 and O157:H7 $\Delta stx_1 stx_2$ serotypes other EHEC serotypes were not affected. Further research is needed to conclude how CBP affect non-O157 EHEC serotypes within mixed ruminal microorganism fluid media.

Table 4.1 Least squares means for growth of enterohemorrhagic *Escherichia coli* (EHEC) O26:H11, O103:H8, O111:H8, O145:H28, O157:H7 and O157:H7 $\Delta stx_1 stx_2$ within bovine mixed rumen microorganism fluid medium, supplemented with 0%, 5%, 10% and 20% powdered citrus by-product (CBP) over time (h)

CBP (%)	Time (h)	O26:H11	O103:H8	O111:H8	O145:H28	O157:H7	O157:H7 $\Delta stx_1 stx_2$
0%							
	0 h	7.55	7.11	6.93	7.55	6.03 ^v	6.46
	2 h	7.68	7.05	6.70	7.58	6.91 ^w	7.00
	4 h	7.60	6.81	6.92	8.06	6.91 ^w	7.19
	6 h	7.73	6.65	7.07	7.83	7.24 ^w	6.98
5%							
	0 h	7.40	7.34 ^y	7.31 ^v	7.57	6.73	6.63
	2 h	7.47	6.86 ^{yz}	7.06 ^v	7.64	6.65	6.41
	4 h	7.43	6.95 ^{yz}	5.85 ^w	7.80	6.77	7.09
	6 h	7.59	6.55 ^z	7.22 ^v	7.80	7.02	6.74
10%							
	0 h	7.43	7.28	6.64	7.73	6.83	6.82 ^{vw}
	2 h	7.43	6.90	7.33	7.56	7.22	6.18 ^w
	4 h	7.36	6.99	7.18	7.64	6.96	7.13 ^v
	6 h	7.66	6.69	7.25	7.63	6.98	4.19 ^x
20%							
	0 h	7.12 ^v	7.13	7.02	7.60	7.03	7.03 ^v
	2 h	7.54 ^v	7.33	6.78	7.52	6.94	6.53 ^v
	4 h	5.96 ^w	7.07	6.65	7.44	7.30	7.34 ^v
	6 h	7.19 ^v	6.66	6.70	7.66	7.00	1.99 ^w

^{v, w, x} Lsmeans within a column, within a treatment, without a common subscript are different between treatment groups if ($P \leq 0.05$)

^{y, z} Lsmeans within a column, within a treatment, without a common subscript within column tend to differ ($P < 0.09$)

Table 4.2 Least squares means for growth of enterohemorrhagic *Escherichia coli* (EHEC) O26:H11, O103:H8, O111:H8, O145:H28, O157:H7 and O157:H7 $\Delta stx_1 stx_2$ within bovine mixed rumen microorganism fluid medium, supplemented with 0%, 5%, 10% and 20% pelleted citrus by-product (CBP) over time (h)

CBP (%)	Time (h)	O26:H11	O103:H8	O111:H8	O145:H28	O157:H7	O157:H7 $\Delta stx_1 stx_2$
0%							
	0 h	9.27 ^x	7.01 ^x	8.47 ^x	9.61	8.48	6.46
	2 h	8.08 ^y	7.01 ^x	9.01 ^{xy}	9.48	8.24	6.28
	4 h	9.45 ^x	8.64 ^y	8.43 ^x	9.67	8.79	6.26
	6 h	9.64 ^x	8.77 ^y	9.48 ^y	9.50	9.01	6.51
5%							
	0 h	9.27 ^{xy}	8.21	8.40 ^{xy}	9.51	9.23 ^x	6.07
	2 h	8.71 ^x	8.26	8.93 ^x	9.60	7.94 ^y	5.92
	4 h	7.29 ^z	7.93	7.98 ^y	9.59	7.96 ^y	6.23
	6 h	9.71 ^y	8.28	8.77 ^x	9.69	8.93 ^x	5.97
10%							
	0 h	9.07 ^{xz}	8.76	8.76 ^{xy}	8.98	8.73 ^{xy}	6.83
	2 h	8.43 ^{xy}	8.50	9.03 ^{xy}	9.01	8.73 ^{xy}	6.94
	4 h	7.90 ^y	8.37	8.66 ^x	9.55	8.40 ^{xy}	6.64
	6 h	9.59 ^z	8.07	9.73 ^y	9.71	9.20 ^y	6.68
20%							
	0 h	9.47 ^x	8.52 ^x	9.03	9.29 ^x	8.00 ^x	6.78
	2 h	8.98 ^x	8.51 ^x	8.92	9.79 ^{xy}	8.95 ^y	6.77
	4 h	7.69 ^y	8.57 ^x	9.09	8.95 ^y	8.71 ^{xy}	6.76
	6 h	9.54 ^x	7.43 ^y	9.46	9.48 ^{xy}	9.05 ^y	7.13

^{x, y, z} Lsmeans within a column, within a treatment, without a common subscript are different between treatment groups if ($P \leq 0.05$)

Literature cited

- Bakkali F, Averbeck S, Averbeck D, Idaomar M. Biological effects of essential oils-a review. *Food and Chem Tox* 2008; 46:446-475.
- Callaway TR, Carroll JA, Arthington JD, Pratt C, Edrington TS, Anderson RC, Galyean ML, Ricke SC, Crandall P, Nisbet DJ. Citrus products decrease growth of *E. coli* O157:H7 and *Salmonella typhimurium* in pure culture and in fermentation with mixed ruminal microorganisms *in vitro*. *Foodborne Path Dis* 2008; 5:621-627.
- Callaway TR, Carroll JA, Arthington JD, Edrington TS, Rossman ML, Carr MA, Krueger MA, Ricke NA, Crandall SC, Nisbet DJ. *Escherichia coli* O157:H7 populations in ruminants can be reduced by orange peel product feeding. *J Food Prot* 2011; 74:1917-1921.
- CDC, 2011a. Preliminary FoodNet Data on the incidence of pathogens transmitted commonly through food-10 states, 2009. Accessed 10 Feb 2012.
<http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5914a2.htm>
- CDC. 2011b. Estimates of Foodborne Illnesses in the United States. Accessed December 23, 2011.
http://www.cdc.gov/nczved/divisions/dfbmd/diseases/ecoli_o157h7/#how_common
- Collier CT, Carroll JA, Callaway TR, Arthington JD. Oral administration of citrus pulp reduces gastrointestinal recovery of orally dosed *Escherichia coli* F18 in weaned pigs. *J Anim Vet Adv* 2010; 9:2140-2045.
- Cotta MA, Russell JB. Effect of peptides and amino acids on efficiency of rumen bacterial protein synthesis in continuous culture. *J Dairy Sci* 1982; 65:226-234.
- Di Pasqua R, Betts G, Hoskins N, Edwards M, Ercolini D, Mauriello G. Membrane toxicity and antimicrobial compounds from essential oils. *J Agric Food Chem* 2007; 55:4863-4870.
- Dusan F, Marian S, Katarina D, Dobroslava B. Essential oils-their antimicrobial activity against *Escherichia coli* and effect on intestinal cell viability. *Toxicol In Vitro* 2006; 20:1435-1445.
- Fisher K, Phillips CA. The effect of lemon, orange and bergamot essential oils and their components on the survival of *Campylobacter jejuni*, *Escherichia coli* O157, *Listeria monocytogenes*, *Bacillus cereus* and *Staphylococcus aureus* *in vitro* and in food systems. *J Appl Microbiol* 2006; 101:1232-1240.

- Free A, Duoss HA, Callaway TR, Carroll JA, Schmidt TB, Donaldson JR. Survival of O157:H7 and non-O157 serotypes of *Escherichia coli* in bovine bile salts and rumen fluid. submitted, 2012.
- Gill AO, Holley RA. Disruption of *Escherichia coli*, *Listeria monocytogenes*, and *Lactobacillus sakei* cellular membranes by plant oil aromatics. *Int J Food Microbiol* 2006; 108:1-9.
- Griffin PM, Tauxe RV. The epidemiology of infections caused by *Escherichia coli* O157:H7, other enterohemorrhagic *E. coli* and the associated hemolytic uremic syndrome. *Epidemiol Rev* 1991; 13:60-98.
- Gustafson JE, Liew YC, Chew S, Markham JK, Bell HC, Wyllie SG, Warmington JR. Effects of tea oil on *Escherichia coli*. *Lett Appl Microbiol* 1998; 26:194-198.
- Kim J, Marshall MR, Wei CI. Antimicrobial activity of some essential oil components against five foodborne pathogens. *J Agric Food Chem* 1995; 43:2839-2845.
- Leyendecker SA, Callaway TR, Anderson RC, Nisbet DJ. Technical note on a much simplified method for collecting ruminal fluid using a nylon paint strainer. *J Sci Food Agric* 2004; 84:387-389.
- Nam IS, Garnsworthy PC, Ahn JH. Supplementation of essential oil extracted from citrus peel to animal feeds decreases microbial activity and aflatoxin contamination without disrupting *in vitro* ruminal fermentation. *Asian-Austral J Anim Sci* 2006; 19:1617-1622.
- Nannapaneni R, Muthaiyan A, Crandall PG, Johnson MG, O'Brien CA, Chalova VI, Callaway TR, Carroll JA, Arthington JD, Nisbet DJ, Ricke SC. Antimicrobial activity of commercial citrus-based natural extracts against *Escherichia coli* O157:H7 isolates and mutant strains. *Foodborne Path Dis* 2008; 5:695-699.
- Oussalah M, Caillet S, Lacroix M. Mechanisms of action of Spanish oregano, Chinese cinnamon, and savory essential oils against cell membranes and walls of *Escherichia coli* O157:H7 and *Listeria monocytogenes*. *J Food Prot* 2006; 69:1046-1055.
- Sambrook J, Russell DW. *Molecular cloning: a laboratory manual*, 3rd edition. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, 2001.
- Stanford K, Bach SJ, Marx TH, Jones S, Hansen JR, Wallins GL, Zahiroddini H, McAllister TA. Monitoring *Escherichia coli* O157:H7 in inoculated and naturally colonized feedlot cattle and their environment. *J Food Prot* 2005; 68:26-33.

USDA-FSIS. FSIS to extend implementation date on routine sampling program for non O157:H7. Accessed 8 Feb 2012.

http://www.fsis.usda.gov/News_&_Events/Const_Update_020812/index.asp

Wells JG, Shipman LD, Greene KD, Sowers EG, Green JH, Cameron DN, Downes FP, Martin ML, Griffin PM, Ostroff SM, Potter ME, Tauxe RV, Wachsmuth IK. Isolation of *Escherichia coli* serotype O157:H7 and other shiga-like-toxin-producing *E. coli* from dairy cattle. J Clin Microbiol 1991; 29:985-989.

CHAPTER V

EVALUATION OF THE COLONIZATION OF *ESCHERICHIA COLI* O157:H7 WITHIN THE SWINE GASTROINTESTINAL TRACT

Abstract

The objective of this study was to monitor the colonization and survival of *Escherichia coli* O157:H7 (ATCC 43888) in the swine gastrointestinal (GI) tract through the use of bioluminescent technology and viable plate counts. Twelve weanling pigs (18-21 d) were orally inoculated with 7.0×10^9 CFU/mL of *E. coli* O157:H7 containing the bioluminescent plasmid pAK1-*lux* at the following intervals: -10 d, -7 d, -2 d and -1 d pre-mortem (0 d). Colonization of *E. coli*+pAK1-*lux* was then analyzed in the duodenum, jejunum, ileum, cecum, colon and rectum by viable plate counts and also through luminescent imaging. The bacterial populations (\log_{10} CFU/mL) increased ($P < 0.001$) as the incubation period decreased, with the 1 d incubation period exhibiting the greatest bacterial populations within all intestinal sections analyzed, with the exception of the jejunum. Following the removal of intestinal contents, each section was individually rinsed to remove any *E. coli* O157:H7+pAK1-*lux* that may have adhered to the intestinal endothelial lining and subsequently analyzed. Data suggest that upon initial exposure to *E. coli* O157:H7, there is a propensity for the bacteria to colonize throughout the GI tract within 1 d. Although the 1 d exposure displayed the greatest populations within the GI rinse, 1 d also displayed similar populations ($P < 0.21$) within all the sections of the 1 d treatment. This study demonstrates that *E. coli*+pAK1-*lux* were successfully monitored

through the GI tract of swine, and a colonization pattern was observed. Further research is needed to validate if *E. coli* O157:H7 transformed with the pAK1-*lux* bioluminescent plasmid could serve as a valid model for long-term *in vivo* studies.

Introduction

In 2011 *E. coli* O157:H7 was responsible for an estimated 1 million pounds of ground beef recalled in the United States due to possible contamination (USDA-FSIS, 2011). Controlling and eliminating fecal shedding of *Escherichia coli* O157:H7 in food animals has been a major issue in pre-harvest food safety and has prompted investigations into intervention strategies to reduce colonization and shedding in livestock (Genovese et al., 2000; Callaway et al., 2003). Many pre-harvest intervention strategies have proven to reduce *E. coli* O157:H7 populations but are not economically feasible to the producer or are not implemented due to the lack of producer utilization and acceptance. Therefore, to date there has been no confirmed method to fully control or eliminate *E. coli* O157:H7 shedding (Huffman, 2002; Callaway et al., 2009).

Historically, swine have not been commonly linked to enterohemorrhagic *E. coli* (EHEC) because unlike cattle, swine are susceptible to EHEC infections (Anderson et al., 2001). *Escherichia coli* O157:H7 is capable of attaching to swine's large intestine epithelial cells in a mechanism similar to that observed in humans (Francis et al., 1986; Tzipori et al., 1986), thus effacing the microvilli and inducing illness (Donnenberg et al., 1993). However, data is surfacing that suggest swine may be asymptomatic carriers of the *E. coli* O157:H7 pathogen (Borie et al., 1997; Anderson et al., 2001; Callaway et al., 2004). For instance, in 1997, 68% of pigs that were sent to harvest in Santiago, Chile, were harboring EHEC (Borie et al., 1997; Anderson et al., 2001). In 1999, *E. coli*

O157:H7 was isolated from swine that were sent to harvest in the Netherlands (Heuvelink et al., 1999). Also in 2007, a nationwide spinach-associated *E. coli* O157:H7 outbreak was linked to feral swine in California (Jay et al., 2007).

Real-time tracking of pathogens can aid in the monitoring and understanding of physiological events, pathogenesis and colonization patterns of bacteria both *in vivo* and *in vitro* (Contag et al., 1995; Contag et al., 1997; Siragusa et al., 1999; Ryan et al., 2010). Successful *in vivo* tracking of *Salmonella typhimurium* transformed with the pAK1-*lux* bioluminescent plasmid was achieved within the swine model (Willard et al., 2002; Moulton et al., 2009b). Moulton et al. (2009b) inoculated swine with *S. typhimurium*+pAK1-*lux* and monitored tracking within the small intestinal tract within a 12 h period. Additionally, *S. typhimurium* and *E. coli* transformed with various *luxCDABE* operons have been successfully implemented within murine gastrointestinal (GI) tract models (Burns-Guydish et al., 2005; Foucault et al., 2010). *Ex vivo* tracking of *E. coli* transformed with the Xen-14 bioluminescent plasmid has been successful in the bovine (Curbelo et al., 2010) and *E. coli* containing pAK1-*lux* within the ovine (Moulton et al., 2009a) reproductive tracts, suggesting that an *in vivo* application of this technology in a swine model is possible. Therefore, the objective of this study was to determine whether *E. coli* O157:H7 transformed with the pAK1-*lux* bioluminescent plasmid could be successfully monitored within the GI tract of weanling pigs.

Materials and Methods

Bacterial Cultivation Conditions

Escherichia coli O157:H7 ATCC 43888 (Δ *stx*₁*stx*₂) was transformed with the pAK1-*lux* plasmid by methods previously described by Moulton et al. (2009a). The *E.*

E. coli O157:H7+pAK1-*lux* is routinely cultured on tryptic soy agar (TSA; Difco Co., Corpus Christi, TX) supplemented with 50 µg/mL ampicillin (AMP) at 37°C. Luminescence was verified after transformation using a Berthold/NightOwl camera equipped with the WinLight 32 software, version 2.51.11901 (Berthold Technologies; Oak Ridge, TN). A fresh, 24 h culture of *E. coli* O157:H7+pAK1-*lux* cultivated in tryptic soy broth (TSB; Difco Co., Corpus Christi, TX) supplemented with 50 µg/mL AMP at 39°C was used for each inoculation challenge.

Swine Trial

All experimental procedures were performed in accordance with the Guide for the Care and Use of Agricultural Animals in Research and Teaching and approved by the Institutional Animal Care and Use Committee of the USDA-ARS. Twelve weanling Yorkshire-Landrace crossbred pigs 18-21 d of age were obtained for this study. Three pigs were orally inoculated via a 10 mL dose oral gavage to achieve an approximate inoculation per pig of 7.0×10^9 CFU/mL at -10 d, -7 d, -2 d, and finally at -1 d prior to euthanization (0 d). Pigs were humanely euthanized and the duodenum, jejunum, ileum, cecum, colon and rectum were aseptically removed from each individual pig; the anterior and posterior ends of each respective section of the GI tract were tied using sterile zip ties.

Gastrointestinal Tract Sections

Each section from each individual weanling pig was stored at 4°C and analyzed within 24 h of collection. Contents from each section were collected, homogenized, and either transferred to a 96-well black sterile plate (Costar®, Corning Inc.; Corning, NY) for imaging or serially diluted in 1X phosphate buffer saline (PBS) and plated onto TSA

supplemented with 50 µg/mL AMP in duplicate for validation of *E. coli*+pAK1-*lux*. A 10 min acquisition period was used for imaging contents from each section using a Berthold/NightOwl camera. Photonic emissions were recorded in RLU, which are measured in photons/pixel second (ph/pix s). Plates were incubated for 12 h at 37°C prior to analysis and were also subsequently imaged to validate luminescence.

Following the collection of contents, the jejunum, ileum, and duodenum were divided into subsections in order to image the entire portion of the tract. The jejunum and ileum were both divided into three sections: posterior, middle and anterior. The duodenum was divided into two sections. Each section or subsection was incubated briefly to reach an internal temperature of 37°C and then subsequently imaged. The RLU values were recorded as previously described above for the intestinal contents analysis.

Gastrointestinal Tract Rinse

Following imaging, each section was stomached (Seward Stomacher® 400 Circular; West Sussex, United Kingdom) at 240 rpm for 1 min in 5 mL of 1X PBS. An aliquot of each dissected section was serially diluted in 1X PBS and plated on TSA supplemented with 50 µg/mL AMP and incubated for 12 h at 37°C. An additional 2 µL aliquot was transferred to a 96-well plate and imaged to measure photonic emissions as described above. After incubation, plates were analyzed and subsequently imaged to validate luminescence.

Statistical Analysis

Data were analyzed as a completely randomized design with repeated measures using PROC MIXED in SAS (SAS Inst. Inc., Version 9.2; Cary, NC). Experimental unit was defined as section, and significance was declared at $P < 0.05$. Pair-wise differences

among least squares means at various sample times were evaluated with the PDIFF statement.

Results

Gastrointestinal Tract Contents

To analyze the colonization of *E. coli* O157:H7 throughout the GI tract of swine, the duodenum, jejunum, ileum, cecum, colon and rectum were collected and contents were removed and analyzed for the presence of *E. coli* O157:H7+pAK1-*lux*. Analysis was based on viable plate count populations in log₁₀ (CFU/mL) within each section, as well as on incubation periods within each section (Table 5.1). The 1 d incubation period had the greatest viable plate count populations observed within the ileum, cecum, colon and rectum. Incubation period ($P < 0.001$) decreased viable plate counts as incubation period increased. There were differences observed within the duodenum ($P < 0.02$), cecum ($P < 0.02$) and rectum ($P < 0.03$) when comparing the 1 d and 2 d incubation periods. There were similar populations observed within 7 d and 10 d populations with the only difference observed within the duodenum ($P < 0.001$), with 10 d displaying decreased populations. The *E. coli*+pAK1-*lux* were initially observed within the jejunum, ileum, cecum and rectum within 1 d incubation period. There were greater populations observed within the jejunum and ileum in the 2 d, 7 d and 10 d incubation periods. There were no differences observed within the colon and rectum populations of the 2 d ($P < 0.90$), 7 d ($P < 0.63$) and 10 d ($P < 0.36$).

Photonic emissions from the *E. coli*+pAK1-*lux* were measured in RLU (ph/pix s) using the Berthold NightOwl camera. The RLU values of the GI content ranged from 38.67 ph/pix s to 53.41 ph/pix s (data not reported in tabular form). The incubation

period ($P < 0.74$) and section ($P < 0.52$) did not affect the capabilities of the pAK1-*lux* plasmid to emit photons, thus exhibiting luminescence. There was no difference in RLU values observed within each individual section; duodenum ($P < 0.27$), jejunum ($P < 0.35$), ileum ($P < 0.19$), cecum ($P < 0.23$), colon ($P < 0.32$) and rectum ($P < 0.12$), respectively.

Gastrointestinal Tract Rinse

After the GI content was removed, each section was stomached in 1X PBS to remove any *E. coli* O157:H7 cells that might have adhered or colonized along the intestinal endothelial lining. The 1 d incubation period had the greatest populations within all rinse of the GI tract and had similar populations when compared to the 2 d incubation period except within the duodenum ($P < 0.003$; Table 5.2). The 7 d and 10 d incubation period data indicated that no difference within populations of the section with the exception of the duodenum ($P < 0.009$), which had greater populations observed at 7 d.

The *E. coli*+pAK1-*lux* initially were observed within the cecum, jejunum and ileum within 1 d and 2 d incubation periods. Greater populations of adhered cells were observed within the cecum, colon and rectum at 1 d incubation period in comparison to the other incubation periods, thus suggesting the possibility of *E. coli* O157:H7 being shed.

The RLU values from the GI sectional rinse were similar to the GI content values. The GI rinse values ranged from 37.09 ph/pix s to 43.09 ph/pix s (not reported in tabular form). The incubation period ($P < 0.61$) or the section ($P < 0.21$) did not affect the photonic emissions or luminescent capabilities of the pAK1-*lux* plasmid. There was no

difference in RLU values observed within each individual section; duodenum ($P < 0.63$), jejunum ($P < 0.27$), ileum ($P < 0.17$), cecum ($P < 0.46$), colon ($P < 0.16$) and rectum ($P < 0.63$).

Discussion

Since most *E. coli* O157:H7 infections are acquired through the consumption of contaminated food products, there have been numerous pre- and post- harvest intervention strategies implemented to reduce prevalence of this dangerous bacterium. Here, we utilized bioluminescent technology to determine pathogen survival and movement within the GI tract of a simple stomached animal. By employing bioluminescence as a monitoring device, researchers can obtain a better understanding of pathogen location and will have potential capabilities of real-time tracking and monitoring of pathogens.

When comparing the GI content treatments over the course of a 1 d infection, significant changes in the colonization pattern of *E. coli* O157:H7 were observed within each section. Overall, there were greater *E. coli* O157:H7 populations observed within the GI rinse than compared to the GI content. This data indicates that the *E. coli* O157:H7 was more apt to adhere to the intestinal endothelial lining rather than remain loose or sloughed within the GI tract.

The greatest populations of *E. coli* O157:H7 within the GI content were observed in the small intestine (duodenum, jejunum, ileum). These data indicate that *E. coli* O157:H7 is able to colonize successfully within the small intestine within 1 d post-exposure. The 1 d incubation period exhibited greater populations that had adhered to the intestinal lining, while the 10 d incubation period exhibited decreased populations within

the cells that did not adhere. This correlates with the increase observed initially within the rectum populations, suggesting that that non-adhered populations are being shed. These data propose that *E. coli* O157:H7 is able to resist bacterial properties within the GI tract such as bile, and still colonize and cause infection. This suggests that upon initial exposure to the pathogen, there is a propensity for the bacteria to colonize throughout the GI tract within 1 d. The *E. coli* O157:H7 populations with an increased incubation period such as 7 d and 10 d post-exposure tend to exhibit decreased populations being colonized and sloughed through the GI tract.

The GI content and GI sectional rinse exhibited similar photonic emission values and results, indicating that each section exhibited relatively the same amount of photonic emissions. All sections exhibited luminescence with little difference observed across incubation groups. Although, the photonic emissions that were obtained were relatively low, these data are in accordance with results previously obtained from our lab, which indicate that photonic emissions from the pAK1-*lux* plasmid decrease over time (Duoss, unpublished). It is possible that the stability of the pAK1-*lux* plasmid is compromised in the absence of AMP *in vivo*, resulting in a decrease of photonic emissions. This is supported by results reported by Moulton et al. (2009a) that indicated in the presence of AMP, there were 100% photon emitting bacteria; and in the absence of AMP there was a significant decrease after 1 d. Although no section exhibited luminescence upon imaging, photonic emissions and viable plate counts ensure that the plasmid, pAK1-*lux* was retained and *E. coli* O157:H7 were present. The use of a more sensitive coupled-camera device, such as the Stanford Photonic Imaging System (XR/MEGA-10Z; Stanford Photonics Inc., Palo Alto, CA), might be more viable for this study, than the NightOwl/Berthold camera used. Therefore, it is still possible that biophotonic

technology could be beneficial to understanding bacterial pathogenesis and colonization patterns *in vivo*. Further research is needed, though, to increase the sensitivity of the procedure used.

Conclusion

Transformation of *E. coli* O157:H7 with the pAK1-*lux* gene cassette did exhibit luminescence within the GI tract implemented within the swine model. Considering that pathogen tracking was successful in the monogastric, simple stomached model, this gives inspiration that pathogen tracking could be utilized within the ruminant, more complex stomached model and be successful within a restricted time frame. Further research is needed to validate if *E. coli* O157:H7 transformed with the pAK1-*lux* bioluminescent plasmid could serve as a valid model for long-term *in vivo* study.

Table 5.1 The least square means of *Escherichia coli* O157:H7+pAK1-*lux* (ATCC 43888) populations (\log_{10} CFU/mL) within contents collected from the gastrointestinal tract of swine

Sections	Log₁₀ CFU/mL						Incubation Period (d)							
	1 d		2 d		7 d		10 d		1 d		2 d		7 d	
	1 d	2 d	2 d	7 d	7 d	10 d	10 d	1 d	2 d	2 d	7 d	7 d	10 d	10 d
Duodenum	6.52 ^a	4.26 ^a	4.40 ^{ab}	4.40 ^{ab}	4.40 ^{ab}	0.12 ^a	0.02	0.01	<0.001	<0.001	0.93	<0.001	<0.001	<0.001
Jejunum	7.73 ^{ab}	8.41 ^b	4.75 ^{be}	4.75 ^{be}	4.75 ^{be}	5.11 ^b	0.41	0.009	0.003	<0.001	<0.001	0.003	0.003	0.66
Ileum	8.80 ^b	8.77 ^b	7.21 ^d	7.21 ^d	7.21 ^d	6.48 ^b	0.97	0.06	0.005	0.07	0.07	0.006	0.006	0.28
Cecum	8.24 ^b	6.17 ^c	3.88 ^{ae}	3.88 ^{ae}	3.88 ^{ae}	4.75 ^{bc}	0.02	<0.001	0.002	0.009	0.009	0.10	0.10	0.32
Colon	6.75 ^a	6.30 ^c	2.97 ^{ac}	2.97 ^{ac}	2.97 ^{ac}	3.59 ^c	0.60	<0.001	0.005	0.003	0.003	0.002	0.002	0.46
Rectum	8.29 ^b	6.40 ^c	2.30 ^c	2.30 ^c	2.30 ^c	3.23 ^c	0.03	<0.001	<0.001	<0.001	<0.001	0.005	0.005	0.27
P-value	0.05	0.03	0.04	0.04	0.04	0.04								

^{a, b, c, d, e} Lsmeans without a common subscript within column differ ($P < 0.05$)

Table 5.2 The least square means of *Escherichia coli* O157:H7+pAK1-*lux* (ATCC 43888) populations (\log_{10} CFU/mL) within rinse obtained from the gastrointestinal tract of swine

Sections	<i>Log</i> ₁₀ CFU/mL									
	Incubation Period (d)									
	1 d		2 d		7 d		10 d		P-values (between incubation periods)	
	1 d	2 d	7 d	10 d	1 d vs 2 d	1 d vs 7 d	1 d vs 10 d	2 d vs 7 d	2 d vs 10 d	7 d vs 10 d
Duodenum	7.72	2.91 ^a	7.34 ^a	4.15 ^x	0.001	0.57	0.005	0.002	0.33	0.009
Jejunum	8.05	7.04 ^b	6.39 ^{ab}	5.82 ^y	0.27	0.08	0.02	0.50	0.21	0.53
Ileum	7.59	7.30 ^b	6.09 ^{ab}	5.38 ^{xy}	0.74	0.11	0.02	0.18	0.04	0.43
Cecum	7.65	7.11 ^b	4.55 ^{bc}	3.52 ^x	0.73	0.06	0.02	0.11	0.03	0.51
Colon	7.23	5.89 ^b	3.46 ^c	5.02 ^{xy}	0.39	0.03	0.16	0.13	0.57	0.32
Rectum	6.34	6.23 ^b	2.75 ^c	3.28 ^{xy}	0.94	0.03	0.17	0.03	0.20	0.35
P-value	0.21	0.03	0.04	0.07						

^{a, b, c} Lsmeans without a common subscript within column differ ($P < 0.05$)
^{x, y} Lsmeans without a common subscript within column tend to differ ($P \leq 0.10$)

Literature cited

- Anderson RC, Callaway TR, Buckley SA, Anderson TJ, Genovese KJ, Sheffield CL, Nisbet DJ. Effects of oral sodium chlorate administration on *Escherichia coli* O157:H7 in the gut of experimentally infected pigs. *Int J Food Microbiol* 2001; 71:125-130.
- Borie C, Monreal Z, Guerrero P, Sanchez ML, Martinez J, Arellano C, Prado V. Prevalence and characterization of enterohemorrhagic *Escherichia coli* isolated from healthy cattle and pigs slaughtered in Santiago, Chile. *Arch Med Vet* 1997; 29:205-212.
- Burns-Guydish SM, Olomu IN, Zhao H, Wong RJ, Stevenson DK, Contag CH. Monitoring age-related susceptibility of young mice to oral *Salmonella enterica* serovar *Typhimurium* infection using *in vivo* murine model. *Pediatr Res* 2005; 58:153-158.
- Callaway TR, Anderson RC, Edrington TS, Elder RO, Genovese KJ, Bischoff KM, Poole TL, Jung YS, Harvey RB, Nisbet DJ. Preslaughter intervention strategies to reduce food-borne pathogens in food animals. *J Anim Sci* 2003; 81(E Suppl 2):E17-E23.
- Callaway TR, Anderson RC, Tellez G, Rosario C, Nava GM, Eslava C, Blanco MA, Quiroz MA, Olguin A, Herradora M, Edrington TS, Genovese KJ, Harvey RB, Nisbet DJ. Prevalence of *Escherichia coli* O157 in cattle and swine in central Mexico. *J Food Prot* 2004; 67:2274-2276.
- Callaway TR, Carr MA, Edrington TS, Anderson RC, Nisbet DJ. Diet, *Escherichia coli* O157:H7, and cattle: A review after 10 years. *Curr Iss Mol Biol* 2009; 11:67-80.
- Contag CH, Contag PR, Mullins JI, Spilman SD, Stevenson DK, Benaron DA. Photonic detection of bacterial pathogens in living hosts. *Mol Microbiol* 1995; 18:593-603.
- Contag CH, Spillman SD, Contag PR, Oshiro M, Eames B, Dennery P, Stevenson DK, Benaron DA. Visualizing gene expression in living mammals using a bioluminescent reporter. *Photochem Pathobiol* 1997; 66:523-531.
- Curbelo J, Moulton K, Willard S. Photonic characteristics and *ex vivo* imaging of *Escherichia coli*-Xen14 within the bovine reproductive tract. *Therio* 2010; 73:48-55.
- Donnenberg MS, Tzipori S, McKee ML, O'Brien AD, Alroy J, Kaper JB. The role of the *eae* gene of enterohemorrhagic *Escherichia coli* in intimate attachment *in vitro* and in a porcine model. *J Clin Invest* 1993; 92:1418-1424.

- Duoss HA. Survival of *Escherichia coli* O157:H7 (ATCC 43888) transformed with either the pAK1-*lux* and pXEN-13 plasmids in mixed bovine rumen and fecal fluid. 2012. Unpublished data.
- Foucault ML, Thomas L, Goussard S, Branchini BR, Grillot-Courvalin C. *In vivo* bioluminescence imaging for the study of intestinal colonization by *Escherichia coli* in mice. *Appl Environ Microbiol* 2010; 76:264-274.
- Francis DH, Collins JE, Duimstra JR. Infection of gnotobiotic pigs with an *Escherichia coli* O157:H7 strain associated with an outbreak of hemorrhagic colitis. *Infect Immun* 1986; 51:953-956.
- Genovese KJ, Anderson RC, Harvey RB, Nisbet DJ. Competitive exclusion treatment reduces the mortality and fecal shedding associated with enterotoxigenic *Escherichia coli* infection in nursery-raised neonatal pigs. *Can J Vet Res* 2000; 64:204-207.
- Heuvelink AE, Zwartkruis-Nahuis JTM, van den Biggerlaar LAM, van Leeuwen WJ. Isolation and characterization of verocytotoxin-producing *Escherichia coli* O157 from slaughter pigs and poultry. *Int J Food Microbiol* 1999; 52:67-75.
- Huffman RD. Current and future technologies for the decontamination of carcasses and fresh meat. *Meat Sci* 2002; 62:285-294.
- Jay MT, Cooley M, Carychao D, Wiscomb GW, Sweitzer RA, Crawford-Miksza L, Farrar JA, Lau DK, O'Connell JA, Millington A, Asmundson RV, Atwill ER, Mandrell RE. *Escherichia coli* O157:H7 in feral swine near spinach fields and cattle, central California coast. *Emerg Infect Dis* 2007; 13:1908-1911.
- Moulton K, Ryan P, Christiansen D, Hopper R, Klausner C, Bennett W, Rodta-Palenik, Willard ST. *Ex vivo* bioluminescence imaging of late gestation ewes following intrauterine inoculation with *lux*-modified *Escherichia coli*. *Comp Immunol Microbiol Infect Dis* 2009a; 32:429-438.
- Moulton K, Ryan P, Lay D, Willard S. Postmortem photonic imaging of *lux*-modified *Salmonella typhimurium* within the gastrointestinal tract of swine following oral inoculation *in vivo*. *J Anim Sci* 2009b; 87: 2239-2244.
- Ryan PL, Christiansen DL, Hopper RM, Walters FK, Moulton K, Curbelo J, Greene JM, Willard ST. Horse Species Symposium: A novel approach to monitoring pathogen progression during uterine and placental infection in the mare using bioluminescence imaging technology and *lux*-modified bacteria. *J Anim Sci* 2010; 89:1541-1551.

- Siragusa GR, Nawotka K, Spilman SD, Contag PR, Contag CH. Real-time monitoring of *Escherichia coli* O157:H7 adherence to beef carcass surface tissues with a bioluminescent reporter. *Appl Environ Microbiol* 1999; 65:1738-1745.
- Tzipori S, Wachsmuth IK, Chapman C, Birner R, Brittingham C, Jackson C, Hogg J. The pathogenesis of hemorrhagic colitis caused by *Escherichia coli* O157:H7 in gnotobiotic pigs. *J Infect Dis* 1986; 154:712-716.
- USDA-FSIS. FSIS Recall Case Archive 2011. Accessed 3 Feb 2012.
http://www.fsis.usda.gov/fsis_recalls/Recall_Case_Archive_2011/index.asp
http://www.fsis.usda.gov/Fsis_Recalls/Open_Federal_Cases/index.asp
- Willard S, Ryan P, Bailey R, Lawrence M, Estill C, Gandy S, Lay JR. Development of a novel paradigm for real-time monitoring of bacterial pathogenicity in swine. *J Anim Sci* 2002; 85(suppl. 1):31. (Abstr).

CHAPTER VI

CONCLUSION

Due to various pre- and post- harvest intervention strategies implemented throughout harvesting and processing plants, the United States food supply is one of the safest in the world. Although safe, foodborne pathogenic bacteria, such as enterohemorrhagic *Escherichia coli*, (EHEC) are still a threat to human health and well-being. Pre-harvest intervention strategies are extensively being researched to discover methods to reduce and/or eliminate EHEC and *E. coli* O157:H7 populations in cattle and decrease the potential for spread and contamination.

Researchers have suggested that the incorporation of citrus by-products within cattle diets may be a successful EHEC and *E. coli* O157:H7 pre-harvest intervention strategy. The essential oils within the citrus by-products possess antimicrobial activities that assert lipophilic properties on pathogens, by increasing permeabilization of the bacteria cell wall and cytoplasm, thus destroying the bacteria. Although previous research suggests that all EHEC function similarly, results presented in this thesis suggest that when citrus by-products in the powder or pelleted variety were supplemented to mixed ruminal microorganism fluid media, EHEC populations were affected differently. While the EHEC serotypes O26:H11, O111:H8, O145:H28 and O157:H7 populations were not affected, the O103:H8 and O157:H7 $\Delta stx_1 stx_2$ (ATCC 43888) had decreased populations. Further research is needed to indicate the effects of citrus by-products on EHEC serotypes and the differences observed within the various EHEC serotypes.

One of the main limitations to pre-harvest food safety intervention strategies is the inability to successfully monitor and track pathogens in real-time. The transformation of *E. coli* O157:H7 with bioluminescent plasmids could aid in the tracking of pathogens. *Escherichia coli* O157:H7 transformed with the bioluminescent plasmids pAK1-*lux* and pXEN-13 grew similarly within mixed ruminal microorganisms fluid media, suggesting that bioluminescent plasmids do not alter the abilities of *E. coli* O157:H7 to thrive in rumen-like conditions of cattle. The implementation of bioluminescent plasmids can also serve as a research tool for comprehending the pathogenesis, pathogen location and physiological systems *in vivo*. *Escherichia coli* O157:H7 transformed with the pAK1-*lux* bioluminescent plasmid was successfully monitored through the gastrointestinal tract of weanling pigs with incubation periods of 10, 7, 2, and 1 day. Successful pathogen tracking within the monogastric, simple stomached model, gives inspiration that pathogen tracking could be utilized within more complex stomached model, such as cattle and be successful within a restricted time frame. Further research is needed to validate if *E. coli* O157:H7 transformed with the bioluminescent plasmids can serve as a valid model for long-term *in vivo* studies.